


RESEARCH ARTICLE

Comparison of diffusing capacity of carbon monoxide (DLCO) and total lung capacity (TLC) between Indigenous Australians and Australian Caucasian adults

Timothy Howarth^{1,2*}, Helmi Ben Saad^{3,4,5}, Ara J. Perez², Charmain B. Atos², Elisha White⁶, Subash S. Heraganahally^{2,6,7}



1 College of Health and Human Sciences, Charles Darwin University, Darwin, Northern Territory, Australia, **2** Darwin Respiratory and Sleep Health, Darwin Private Hospital, Darwin, Northern Territory, Australia, **3** Université de Sousse, Faculté de Médecine de Sousse, Laboratoire de Physiologie, Sousse, Tunisia, **4** Department of Physiology and Functional Exploration, Farhat HACHED Hospital, Sousse, Tunisia, **5** Heart Failure Research Laboratory (LR12SP09), Farhat HACHED Hospital, Sousse, Tunisia, **6** Department of Respiratory and Sleep Medicine, Royal Darwin Hospital, Darwin, Northern Territory, Australia, **7** College of Medicine and Public Health and Northern Territory Medical Programme, Flinders University, Adelaide, South Australia, Australia

* timothypaul.howarth@cdu.edu.au

OPEN ACCESS

Citation: Howarth T, Saad HB, Perez AJ, Atos CB, White E, Heraganahally SS (2021) Comparison of diffusing capacity of carbon monoxide (DLCO) and total lung capacity (TLC) between Indigenous Australians and Australian Caucasian adults. PLoS ONE 16(4): e0248900. <https://doi.org/10.1371/journal.pone.0248900>

Editor: Koustubh Panda, University of Calcutta, INDIA

Received: November 24, 2020

Accepted: March 7, 2021

Published: April 2, 2021

Copyright: © 2021 Howarth et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All files are available from the Charles Darwin University eSpace Institutional Repository database (DOI: [10.25913/4m2r-ek13](https://doi.org/10.25913/4m2r-ek13)).

Funding: SSH as director of Darwin Respiratory and Sleep Health received a voluntary donation from Boehringer Ingelheim (<https://www.boehringer-ingelheim.com.au/>) which as such did not attract any grant number. The funders had no role in study design, data collection and analysis,

Abstract

Background and objective

Currently there is paucity of evidence in the literature in relation to normative values for diffusing capacity of carbon monoxide (DLCO) and total lung capacity (TLC) among Indigenous Australians. Hence, in this study we assessed the DLCO and TLC parameters among Indigenous Australians in comparison to Australian Caucasian counterparts.

Methods

DLCO and TLC values were assessed and compared between Indigenous Australians and Australian Caucasians matched for age, sex and body mass index, with normal chest radiology.

Results

Of the 1350 and 5634 pulmonary function tests assessed in Indigenous Australian and Australian Caucasian adults respectively, a total of 129 Indigenous Australians and 197 Australian Caucasians met the inclusion criteria. Absolute DLCO and TLC values for Indigenous Australians were a mean 4.3 ml/min/mmHg (95% CI 2.86, 5.74) and 1.03 L (95% CI 0.78, 1.27) lower than Australian Caucasians ($p < 0.01$). Percentage predicted values were 15.38 (95% CI 11.59, 19.17) and 16.63 (95% CI 13.59, 19.68) points lower for DLCO and TLC, respectively. Lower limit of normal (LLN) values did not significantly differ between groups, however a significantly greater proportion of Indigenous Australians recorded values below the LLN in comparison to Australian Caucasians for DLCO (64 vs. 25%, $p < 0.01$) and TLC (66 vs. 21%, $p < 0.01$). Significant differences for the interaction of sex on DLCO and TLC

decision to publish, or preparation of the manuscript.

Competing interests: We also want to declare that this study was partially/voluntarily supported by Boehringer Ingelheim pharmaceutical company to support respiratory research in Aboriginal Australian population of the Northern Territory of Australia. Boehringer Ingelheim did not have any role in the study design, data collection, analysis, or interpretation. The funds were utilised to employ research assistants and statisticians for this study. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

Abbreviations: CI, Confidence Interval; DLCO, Diffusion capacity of carbon monoxide; FEV₁, Forced Expiratory Volume in one second; FVC, Forced Vital Capacity; GLI, Global Lung function Initiative; HREC, Human Research Ethics Committee; LFPs, Lung function parameters; LLN, Lower limit of normal; NT, Northern Territory; PFT, Pulmonary function test; TEHS, Top End Health Service; TLC, Total lung capacity.

were noted in Australian Caucasians, with reduced or absent sex differentiation among Indigenous Australians.

Conclusions

There are significant differences in DLCO and TLC parameters between Indigenous Australian compared to Australian Caucasians. Appropriate DLCO and TLC norms need to be established for Indigenous Australians.

Introduction

Indigenous Australians experience a significantly higher burden of respiratory disease in comparison to their non-Indigenous peers [1–4]. Multiple factors underpin this high burden of respiratory disease in this population, from higher rates of premature birth and low birth weight [5], to childhood infections [6] and higher rates of smoking [7, 8]. Currently there is scarcity of published data for adult reference values of normative lung function parameters (LFPs) among Indigenous Australians [9, 10]. Recent work has sought to address this gap in the literature with reports demonstrating significantly reduced Forced Vital Capacity (FVC) and Forced Expiratory Volume in one second (FEV₁), yet typically nearly preserved FEV₁/FVC ratio [11–13]. Moreover, a recent study from our centre found the Global Lung function Initiative (GLI-2012) norms do not fit for Indigenous Australian adults, regardless of which ethnicity option was selected, including “others/mixed” [14]. Normative reference values for diffusing capacity of carbon monoxide (DLCO) and total lung capacity (TLC) have also yet to be established within the Indigenous Australian population [15].

DLCO, a measure of alveolar surface area and diffusion capacity, appears to be a more sensitive measure of morbidity in comparison to FVC or FEV₁ [16]. FVC and/or FEV₁ may remain relatively stable over time even in the presence of advancing pulmonary disease and worsening respiratory symptoms, while DLCO could demonstrate significant decline/abnormality [17, 18]. Furthermore, DLCO has a significantly greater correlation to radiographic indices of pulmonary disease than any other pulmonary function test (PFT) parameters [19]. This factor may be particularly important in resource poor settings, such as those experienced by many Indigenous Australians as access to radiography is extremely limited among patients living in regional and remote communities. DLCO (expressed as a percentage of normative reference values) is among the preoperative tests performed before undergoing pulmonary surgical interventions (**eg**, lobectomy up to a pneumonectomy). In practice, any patient free from cardiovascular risk factors, with a DLCO >80% (in addition to a FEV₁ >80%) can undergo surgical pulmonary interventions, such as pneumonectomy without further investigation—thus it is critical normative reference values actually reflect normative values within the population being tested [20].

In a similar vein, accurate reference values of TLC have at least three factors of primary care significance. First, it permits the determination of the TLC limits of normal (**eg**: lower limit of normal (LLN)), and therefore facilitates accurate diagnosis of specific respiratory disease patterns (**ie**: restrictive ventilatory defect (TLC < LLN) [21] and lung hyperinflation (TLC > upper limit of normal) [22]. Second, TLC allows more meaningful interpretations of FEV₁ and FVC [23]. Third, reduced TLC (**eg**: TLC < LLN) correlates with both morbidity & mortality, quality of life and physical activity [24, 25]. Variations in TLC may be a function of intrinsic factors (body stature and body composition distribution) within the Indigenous

Australian population. However, significant extrinsic barriers also co-exist throughout the life course in this population, such as maternal smoking [26], lower birthweight and gestational age [27], infections through infancy and childhood [6] and higher adolescent and adulthood smoking rates [28], which may contribute to impaired lung growth and development, translating to reduced adult TLC values. Hence, it may be reasonable to document the normative DLCO and TLC parameters in this adult Indigenous Australian population. Therefore, the aim of this study was to evaluate the DLCO and TLC values among radiographically healthy Indigenous Australians and compare these to Australian Caucasians residing in the Top End Health Service (TEHS) region of the Northern Territory (NT) of Australia.

Methods

Setting and ethical approval

This retrospective study was conducted at the Respiratory and Sleep service based at the Royal Darwin Hospital and Darwin Respiratory and Sleep Health, Darwin Private Hospital in the TEHS region, NT of Australia. According to the Australian Bureau of Statistics, about 3.3% of the Australian population identify as Indigenous Australians [29, 30]. In the NT, about 30% of the population identify as Indigenous Australian, the highest proportion amongst all Australian states and territories. This study was approved by the Human Research Ethics Committee of the NT, TEHS and Menzies School of Health Research. (Reference no: HREC 2019–3445). The HREC advised specific written/oral consent was not required as data was routinely collected and analysed anonymously.

Study participants

Study participants included were Indigenous Australian and Australian Caucasian adults living in the TEHS region, who underwent a pulmonary function test between 2012 and 2020 and were referred for PFTs by a primary health practitioner, respiratory specialist or other specialist physician as a part of routine clinical care. A large part of the methodology of this study has been described in our earlier report [14].

Australian Caucasian subject

During the study period, 5634 PFTs were performed on self-identified Australian Caucasian adults. For comparative analysis, subjects who had normal chest radiology were included. The original database was searched for Australian Caucasians to be matched as close as possible for sex, age ranges (18–25, 25–35, 35–55 & 55+ years) and corpulence status (underweight (body mass index (BMI) ≤ 18.5 kg/m²), normal weight (BMI 18.6–24.9 kg/m²), overweight (BMI 25.0–29.9 kg/m²), obese (BMI ≥ 30.0 kg/m²)) against Indigenous Australian counterparts.

PFTs

All PFTs including DLCO and TLC were performed according to the American thoracic and the European respiratory societies guidelines/recommendations, including calibration of equipment and quality assurance [16, 31] and as detailed in our previous reports [11, 14]. For measuring TLC and DLCO, the portable single-breath diffusing capacity device (EasyOne Pro[®], ndd Medical Technologies) was used [32, 33]. Only PFTs graded as acceptable for session quality were included in this study. For acceptable quality for DLCO testing inspiratory volume $>85\%$ best vital capacity was used. TLC was measured by single breath manoeuvre technique [34]. Limitations to single-breath TLC measurement in the presence of airway obstruction are corrected for in EasyOne Pro device using Punjabi et al.'s equation [35]. In our

centre and in the absence of specific respiratory reference values for the Indigenous Australian population, the predicted normative values for TLC and DLCO were calculated using the National Health and Nutrition Examination Survey Caucasian reference set's (no ethnic correction was used) [36]. The following parameters were determined: TLC sb (single breath) (L, %), and DLCO (ml/min/mmHg, %). For the purpose of this study only DLCO and TLC were included in the analysis. No correctional factors for haemoglobin were applied, hence DLCO adj (adjusted) values and carbon monoxide transfer coefficient (= DLCO/alveolar volume) were not included in the analysis.

Clinical data collection and study subject inclusion/exclusion criteria

As per standard protocol, participants age (on the date of testing), sex, height and weight were recorded. Smoking history was recorded to identify current-, past- or never- smokers, and to quantify the pack years of smoking [37]. Participants who underwent PFTs and had chest radiology (Chest X-Ray and or computed tomography chest) performed within a calendar year of the PFT date were initially included in the study, to assess the presence/absence of any radiological pulmonary abnormalities. Only participants with normal chest radiology were included for analysis.

Statistical methods

Continuous parameters were tested for normality with the Shapiro Wilks distribution test. Age, height, weight, BMI (weight/height²), DLCO and TLC were found to deviate only slightly from normal distribution, however smoking pack years deviated to a significant extent and thus was treated as non-parametric. Non-parametric parameters were presented as medians (interquartile ranges), normally distributed parameters as means (95% confidence intervals (CIs)), and categorical parameters as numbers (%). Demographic parameters were compared between Indigenous Australian and Australian Caucasian adults with two-tailed proportions z-tests for categorical parameters, two-tailed students t-test for parameters approximating normal distribution and equality of medians test for the non-parametric. DLCO and TLC absolute and predicted values were tested against each other by age (≤ 35 & > 35 years), sex and Indigenous status using unpaired, two-sided, students t-tests. The proportion of participants outside of norms (ie, $< LLN$) for DLCO and TLC were tested by age, sex and Indigenous status using two-tailed proportions z-tests. Kernel density plots utilising Epanechnikov kernels were produced for each age (≤ 35 & > 35 years), sex (female & male) and Indigenous strata to demonstrate differences in distribution of DLCO and TLC. Multivariate linear regression models were developed to define the effect of known significant factors (age, sex, height, weight and smoking status) alongside Indigenous status on absolute DLCO and TLC, reporting beta coefficients (95% CIs), factor p-values, model p-values and adjusted Pearson's determination coefficient " r^2 ". All data were analysed in STATA IC 15 (StataCorp, Texas) and alpha was set to 0.05 throughout.

Results

Baseline clinical parameters

A total of 326 subjects (129 Indigenous Australian (80 females) and 197 Australian Caucasians (135 females)) were included. Indigenous Australian participants were significantly younger than Australian Caucasians ($p = 0.01$) (Table 1). BMI did not significantly differ between the two populations ($p = 0.96$), but a significantly higher proportion of Indigenous Australians were in the underweight category (7 vs. 1%, $p < 0.01$) and a lower though not significantly

Table 1. Demographics and anthropometrics of Indigenous Australian and Australian Caucasians participants (n = 326).

		Indigenous Australian n = 129	Australian Caucasians n = 197	p-value
Sex (female)		80 (62%)	135 (69%)	0.23
Age (years)		46.22 (44.05, 48.39)	49.64 (48.07, 51.21)	0.01*
Age ranges (years)	< 25	8 (6%)	1 (1%)	<0.01*
	25 ≤ 35	20 (16%)	21 (11%)	0.27
	35 ≤ 55	66 (51%)	104 (53%)	0.88
	> 55	35 (27%)	71 (36%)	0.067
Height (m)		1.66 (1.64, 1.67)	1.68 (1.67, 1.69)	0.044*
Weight (kg)		84.08 (80, 88.16)	86.1 (83.19, 89)	0.41
Body mass index (BMI, kg/m ²)		30.51 (29.09, 31.93)	30.47 (29.53, 31.4)	0.96
Corpulence status	Underweight	9 (7%)	2 (1%)	<0.01*
	Normal weight	24 (19%)	30 (15%)	0.42
	Overweight	32 (25%)	68 (35%)	0.063
	Obese	64 (50%)	97 (49%)	0.95
Smoking status	Non-smoker	23 (18%)	67 (34%)	<0.01*
	Former smoker	34 (27%)	87 (44%)	<0.01*
	Current smoker	70 (55%)	43 (22%)	<0.01*
Pack years †		13 (4, 26)	15 (5.3, 30)	0.77

Data were expressed as mean (95% confidence interval) and number (%).

Differences between groups tested with two-tailed students T-test for continuous parameters and proportions z-test for categorical parameters.

†Pack years data were available for 97 & 118 Indigenous Australian and Australian Caucasian subjects, respectively.

†This variable was non-parametrically distributed, thus displayed as median (interquartile range) and tested via equality of medians test.

*Denote statistically significant differences (p<0.05).

<https://doi.org/10.1371/journal.pone.0248900.t001>

proportion in the overweight category (25 vs. 35%, p = 0.063). A significantly higher proportion of Indigenous Australians reported current smoking (55 vs. 22%, p<0.01), however reported pack years did not differ by Indigenous status (13 (4, 26) vs. 15 (5, 30), p = 0.77).

DLCO data

Indigenous Australian participants displayed significantly reduced absolute DLCO values, with a mean difference of 4.3 ml/min/mmHg (2.86, 5.74, p<0.01) (Table 2). Percentage predicted values were a mean 15.38 (11.59, 19.17, p<0.01) percentage points lower. After being stratified by age and sex, differences were noted in the effect of Indigenous status between strata. Where for females the mean differences by Indigenous status were 11.44 (-2.17, 25.05) and 13.95 (9.15, 18.75) percentage points for under and over 35 years respectively, for males they were 21.33 (6.1, 36.56) and 21.33 (13.45, 29.23).

Absolute LLN values for DLCO did not significantly differ across the total sample (p = 0.99), however were significantly reduced among Indigenous Australian females aged ≤ 35 years (p<0.01) and Indigenous Australian males aged > 35 years (p<0.01). However, a significantly greater proportion of Indigenous Australians both overall, and in each stratum aside from females ≤ 35 years, recorded DLCO values below the LLN (64 vs. 25%, p<0.01).

The distribution of absolute DLCO scores followed a similar pattern for Indigenous Australian and Australian Caucasians for females (any age) and for males > 35 years, though displaced to lower values for Indigenous Australians. However, for males ≤ 35 years the distribution was flipped (Fig 1). Where Australian Caucasian participants displayed an

Table 2. Diffusing capacity of carbon monoxide (DLCO) values for participants by Indigenous status, stratified by sex and age (≤ 35 & > 35 years).

	Indigenous Australian (n = 129)	Australian Caucasians (n = 197)	p-value
Total sample			
DLCO (ml/min/mm/Hg)	18.75 (17.65, 19.85)	23.05 (22.13, 23.97)	$<0.01^*$
DLCO (%)	71 (68, 74)	86 (84, 89)	$<0.01^*$
DLCO LLN	20.16 (19.45, 20.86)	20.4 (19.82, 20.99)	0.99
DLCO < LLN	82 (64%)	49 (25%)	$<0.01^*$
Females			
≤ 35 years	n = 14 (18%)	n = 11 (9%)	
DLCO (ml/min/mm/Hg)	20.23 (17.59, 22.87)	25.87 (21.79, 29.95)	$<0.01^*$
DLCO (%)	78.29 (70.27, 86.3)	89.73 (76.93, 102.53)	0.10
DLCO LLN	20.19 (18.76, 21.63)	23.03 (22.07, 23.99)	$<0.01^*$
DLCO < LLN	7 (50%)	4 (36%)	0.50
> 35 years	n = 66 (82%)	n = 124 (91%)	
DLCO (ml/min/mm/Hg)	16.47 (15.32, 17.61)	20.03 (19.3, 20.76)	$<0.01^*$
DLCO (%)	70.21 (66.05, 74.38)	84.16 (81.42, 86.9)	$<0.01^*$
DLCO LLN	17.7 (17.05, 18.35)	18.1 (17.68, 18.53)	0.29
DLCO < LLN	42 (64%)	32 (26%)	$<0.01^*$
Males			
≤ 35 years	n = 14 (29%)	n = 11 (18%)	
DLCO (ml/min/mm/Hg)	26.59 (22.13, 31.05)	33.72 (29.59, 37.84)	0.02*
DLCO (%)	77.21 (64.83, 89.6)	98.55 (89.79, 107.3)	0.01*
DLCO LLN	27.5 (26.3, 28.7)	27.14 (24.61, 29.66)	0.76
DLCO < LLN	9 (64%)	1 (9%)	0.01*
> 35 years	n = 35 (71%)	n = 51 (82%)	
DLCO (ml/min/mm/Hg)	19.33 (17.1, 21.57)	27.48 (25.64, 29.32)	$<0.01^*$
DLCO (%)	67 (60.08, 73.92)	88.33 (83.68, 92.99)	$<0.01^*$
DLCO LLN	21.83 (20.92, 22.74)	23.98 (22.97, 24.98)	$<0.01^*$
DLCO < LLN	24 (69%)	12 (24%)	$<0.01^*$

Abbreviations: DLCO, diffusing capacity of carbon monoxide; LLN, lower limit of normal.

Data were expressed as mean (95% confidence interval) and number (%).

DLCO (%), LLN) data were derived from the NHANES (National Health and Nutrition Examination Survey) norms.

Differences between groups tested with two-tailed students T-test for continuous parameters and proportions z-test for categorical parameters.

*Denote statistically significant differences ($p < 0.05$).

<https://doi.org/10.1371/journal.pone.0248900.t002>

extended tail toward low DLCO values, Indigenous Australians displayed a tail toward high DLCO values. Furthermore, the disparity in DLCO values by sex was significantly reduced among Indigenous Australians, particularly in those aged > 35 years.

In the development of the multivariate regression model, a significant interaction was noted between Indigenous status and sex, thus separate models were developed for Indigenous Australian and Australian Caucasians (Table 3). Significant differences are seen between the two models for multiple factors. Sex did not display any significant association within Indigenous Australian subjects ($p = 0.47$) whereas it did for Australian Caucasians ($p < 0.01$). Former and current smoking did not reach significance for effect on DLCO values for Indigenous Australians, though current smoking did significantly negatively correlate in Australian Caucasians ($p < 0.01$). The effect of advancing age differed notably between groups, as exhibited by the difference in confidence interval to mean by Indigenous status, with a larger effect visible in the Australian Caucasian group (Fig 2).




Fig 1. Kernel density plots of absolute diffusing capacity of carbon monoxide (DLCO) values by age group and sex. Bandwidth set at 3.00, Epanechnikov kernels.

<https://doi.org/10.1371/journal.pone.0248900.g001>

Table 3. Multivariate regression on diffusing capacity of carbon monoxide (DLCO) value outcome reporting coefficients (95% CIs).

	Indigenous Australian		Australian Caucasians	
	Beta (95% CIs)	p-value	Beta (95% CIs)	p-value
Sex (Female reference)	0.79 (-1.35, 2.92)	0.466	5.06 (3.32, 6.79)	<0.01
Age (years)	-0.16 (-0.22, -0.09)	<0.001	-0.21 (-0.27, -0.15)	<0.01
Height (cm)	0.21 (0.09, 0.35)	0.001	0.15 (0.06, 0.25)	<0.01
Weight (kg)	0.1 (0.06, 0.14)	<0.001	0.05 (0.02, 0.08)	<0.01
Former smoker (non-smoker reference)	-2.21 (-4.6, 0.19)	0.071	-0.23 (-1.66, 1.2)	0.75
Current smoker (non-smoker reference)	-2.04 (-4.16, 0.08)	0.059	-3.21 (-4.95, -1.46)	<0.01
Model R² & p-value	0.515	<0.001	0.584	<0.01

Data were expressed as means (95% confidence interval).

R²: adjusted form of Pearson's product moment correlation coefficient.

Values fit within the 'Moderate' correlation range (0.5–0.7).

<https://doi.org/10.1371/journal.pone.0248900.t003>




Fig 2. Coefficients plot of multivariate regression effects on diffusing capacity of carbon monoxide (DLCO).

<https://doi.org/10.1371/journal.pone.0248900.g002>

TLC data

Indigenous Australian study participants displayed significantly reduced absolute TLC, with a mean difference of 1.03 L (0.78, 1.27, $p < 0.01$) between groups (Table 4). Percentage predicted values were a mean 16.63 (11.59, 19.68, $p < 0.01$) percentage points lower. When stratified by age and sex, differences were noted in the effect of Indigenous status between strata. Where for females the mean difference by Indigenous status increased with age from 8.88 (-2.08, 19.83) to 17.99 (13.71, 22.27) percentage points for under and over 35 years respectively, for males the mean difference decreased from 20.1 (9.56, 30.64) to 15.45 (10.0, 20.9).

Absolute LLN values for TLC did not significantly differ across the total sample ($p = 0.35$), nor any strata aside from males aged > 35 years ($p < 0.01$). However, a significantly greater proportion of Indigenous Australians both overall, and in each stratum aside from females ≤ 35 years, recorded TLC values below the LLN (66 vs. 21%, $p < 0.01$).

The distribution of absolute TLC scores followed a similar pattern for Indigenous Australian and Australian Caucasians females and males (any age), though displaced to lower values for Indigenous Australians (Fig 3). The disparity in TLC values by sex was significantly reduced among Indigenous Australians compared to Australian Caucasians.

Significant differences were noted between the Indigenous Australian and Australian Caucasians in the multivariate regression models for multiple factors (Table 5). Sex bordered on significance for Indigenous Australians ($p = 0.057$), while it was strongly significant for Australian Caucasians ($p < 0.01$). Weight however, displayed a significant association in the Indigenous Australian model ($p = 0.05$), and no association in the Australian Caucasian model ($p = 0.94$) (Fig 4). The overall fit of the models also showed significant difference, accounting for 34.7% of variance in outcomes among Indigenous Australians, and 58% of variance in Australian Caucasians.

Discussion

This study demonstrated that Australian Indigenous people have significantly lower DLCO and TLC values for both absolute and percentage predicted values in both sexes and across ages in comparison to their Australian Caucasian counterparts. However, LLN values for DLCO and TLC did not demonstrate significant differences between the groups.

Table 4. Total Lung Capacity (TLC) values for participants by Indigenous status and stratified by sex and age (≤ 35 & > 35 years).

	Indigenous Australian (n = 129)	Australian Caucasians (n = 197)	p-value
Total sample			
TLC (L)	4.06 (3.87, 4.25)	5.09 (4.93, 5.24)	<0.01*
TLC (%)	73.31 (70.62, 76)	89.94 (88.2, 91.69)	<0.01*
TLC LLN	4.51 (4.35, 4.67)	4.62 (4.48, 4.76)	0.35
TLC < LLN	85 (66%)	41 (21%)	<0.01*
Females			
≤ 35 years	n = 14 (18%)	n = 11 (9%)	
TLC (L)	4.09 (3.55, 4.63)	4.81 (4.31, 5.31)	0.048*
TLC (%)	81.21 (72.5, 89.92)	90.09 (83.32, 96.87)	0.11
TLC LLN	4.04 (3.76, 4.31)	4.33 (4.08, 4.58)	0.11
TLC < LLN	6 (43%)	2 (18%)	0.19
> 35 years	n = 66 (82%)	n = 124 (91%)	
TLC (L)	3.59 (3.38, 3.81)	4.57 (4.44, 4.7)	<0.01*
TLC (%)	73.14 (69.18, 77.1)	91.13 (88.79, 93.46)	<0.01*
TLC LLN	3.95 (3.8, 4.09)	4.01 (3.94, 4.09)	0.36
TLC < LLN	43 (65%)	18 (15%)	<0.01*
Males			
≤ 35 years	n = 14 (29%)	n = 11 (18%)	
TLC (L)	4.97 (4.37, 5.57)	6.44 (5.78, 7.1)	<0.01*
TLC (%)	71.36 (63.43, 79.28)	91.45 (84, 98.91)	<0.01*
TLC LLN	5.78 (5.48, 6.07)	5.88 (5.53, 6.23)	0.62
TLC < LLN	11 (79%)	2 (18%)	<0.01*
> 35 years	n = 35 (71%)	n = 51 (82%)	
TLC (L)	4.57 (4.21, 4.93)	6.11 (5.82, 6.4)	<0.01*
TLC (%)	71.26 (66.28, 76.23)	86.71 (83.63, 89.78)	<0.01*
TLC LLN	5.27 (5.06, 5.48)	5.87 (5.66, 6.08)	<0.01*
TLC < LLN	25 (71%)	19 (37%)	<0.01*

Abbreviations: TLC, Total lung capacity; LLN, Lower limit of normal.

Data were expressed as mean (95% confidence interval) and number (%).

TLC (%), LLN) data were derived from the NHANES (National Health and Nutrition Examination Survey) norms.

Differences between groups tested with two-tailed students T-test for continuous parameters and proportions z-test for categorical parameters.

*Denote statistically significant differences ($p < 0.05$).

<https://doi.org/10.1371/journal.pone.0248900.t004>

Recently, it has been recognised that ethnically diverse normative data for lung diffusing capacity and lung volumes needs to be a research priority for various ethnic populations around the world [38]. To the authors' best knowledge, this is the first study to document LFPs for DLCO and TLC among Indigenous Australian people in comparison to the Australian Caucasian population, especially from the TEHS, NT region of Australia, where 30% of the population is of Indigenous Australian descent compared to the total 3.3% across Australia as a whole.

There have been only a few studies in the past illustrating the spirometry patterns among the Indigenous Australian population [39–43]. Some studies have demonstrated FEV₁ and FVC values to be 20 to 30% lower [43], yet with a nearly preserved FEV₁/FVC ratio [11, 14] among the Indigenous Australian population in comparison to non-Indigenous Australians. One previous study in the Australian setting has compared normative references for DLCO among middle aged Australians with other reference equations [44]. However, none of the




Fig 3. Kernel density plots of absolute Total Lung Capacity (TLC) values by age group and sex. Bandwidth set at 0.400, Epanechnikov kernels.

<https://doi.org/10.1371/journal.pone.0248900.g003>

Table 5.

	Indigenous Australian		Australian Caucasian	
	Beta (95% CIs)	p-value	Beta (95% CIs)	p-value
Sex (Female reference)	0.41 (-0.01, 0.83)	0.057	0.76 (0.46, 1.06)	<0.01
Age (years)	-0.01 (-0.02, 0)	0.108	0 (-0.01, 0.01)	0.60
Height (cm)	0.05 (0.02, 0.07)	0.001	0.06 (0.05, 0.08)	<0.01
Weight (kg)	0.01 (0, 0.01)	0.050	0 (-0.01, 0.01)	0.94
Former smoker (non-smoker reference)	-0.2 (-0.67, 0.27)	0.401	0.1 (-0.14, 0.35)	0.41
Current smoker (non-smoker reference)	-0.07 (-0.49, 0.34)	0.726	-0.09 (-0.39, 0.21)	0.56
Model R² & p-value	0.347	<0.001	0.580	<0.01

Data were expressed as means (95% confidence interval).

R²: adjusted form of Pearson's product moment correlation coefficient.

Values fit within the 'Mild' (0.3–0.5) or 'Moderate' (0.5–0.7) correlation range.

<https://doi.org/10.1371/journal.pone.0248900.t005>




Fig 4. Coefficients plot of multivariate regression effects on Total Lung Capacity (TLC).

<https://doi.org/10.1371/journal.pone.0248900.g004>

aforementioned studies have explored the DLCO and TLC parameters among Indigenous Australian people in comparison to the Australian Caucasian cohort, including the recently published GLI normative data among Aboriginal or Torres Strait Islander children and young adults [45]. Hence, we believe that our study is of significant value to better understand the LFPs in the Indigenous Australian population, particularly in relation to DLCO and TLC parameters. Moreover, the GLI studies [46, 47] including “the GLI-2012 contemporary adult Australian spirometry study” [10], do not contain adequate representation of the Indigenous Australian people. Currently there are no well-established spirometry normative reference equations for the adult Indigenous Australian population and a recent study including children and adults with a smaller samples size has shown the GLI-2012 reference equation do not fit FEV₁ or FVC for any other ethnic groups [48]. Similarly, our recently published study involving older adults demonstrated that the GLI-2012 norms do not fit for Australian Indigenous adults regardless of which ethnicity option was selected, including “others/mixed” [14].

The mechanisms underpinning the lower DLCO and TLC values observed in Indigenous cohort in this study in comparison to the Caucasian Australians are not entirely clear [13]. However, it may be reasonable to presume several genetic/intrinsic or extrinsic/environmental factors may be influential for such observation. These factors may include maternal smoking rates [26], lower birthweight [5, 27], increased childhood infections [6] and adolescent smoking [7, 8, 28]—aspects which have previously been observed to impact upon the applicability of normative values for a population [13, 49]. It is worth noting that Caucasian adults have typically shown significantly larger values for FVC compared to multiple different ethnic populations indicating the potential effect of evolutionary factors—thus the development of the GLI-2012 standard norms for multiple ethnicities [46]. Larger FVC values imply larger TLC values, though the latter is not always measured.

The prevalence of respiratory disorders is increasing worldwide, and in Australia, this is indeed true for the Indigenous population, imposing a significant burden on health care utilisation and cost [1–6, 11–14]. LFPs are crucial and integral in the diagnosis, management and monitoring of several respiratory disorders. Hence, it may be meaningful to make efforts to document normative LFPs not only in our Indigenous Australian people as in this study, but also in other underprivileged and vulnerable First Nations Indigenous populations around the

world. The present study results may be of value to compare the findings to other ethnic groups and also to explore the fit to other ethnic Indigenous population. Moreover, in the absence of any comparative normative reference equation for DLCO and TLC in this reference population, the results of this study may indicate that LLN values may be a better indicator for clinical decision making in the Indigenous Australian population. However, further studies are warranted.

Limitations

The results of this study must be viewed with caution. Our findings are applicable to the Indigenous Australian people living in the Top End NT of Australia and may not represent other Indigenous Australians living in other states and territory or other First Nations Indigenous people around the world. Moreover, the DLCO and TLC results represented in this study were performed on portable DLCO ndd device [33, 34, 50]. It is not clear if the results, particularly for TLC, would differ if the tests were performed using a body plethysmography device. However, measurement of DLCO and TLC values via body plethysmography would face significant challenges in this population, where 81% of Indigenous people reside in remote or very remote communities, many of which may be accessed only seasonally by light aircraft (S1 and S2 Figs) within a vast geographic area of 245,000 km² (94,595.0 sq mi) and a population density of 0.16 people per kilometre, making such a study virtually infeasible [29, 30]. Therefore, the use of a portable DLCO ndd device is the most reasonable method to assess LFP's for Indigenous subjects during respiratory specialist outreach visits to remote Indigenous communities (S3 Fig)—a unique model adopted for respiratory care in the TEHS region of NT of Australia [2, 3, 11]. The authors are also very aware and acknowledge that the sample size may be less than ideal for this study. However, given the fact that Australian Indigenous people have higher incidence of smoking [7, 8, 28, 45], are known to have a higher proportion of respiratory disorders [1–6] and experience geographic isolation, unprecedented challenges are imposed in recruiting study participants without pre-existing lung disease or smoking history for a study such as this.

Conclusion

Despite some limitations and challenges, this is the first study to document the DLCO and TLC among Indigenous Australian people and compare it to Australian Caucasians. Indigenous Australians displayed significantly lower DLCO and TLC values for both absolute and percentage predicted in both sexes and across age ranges in comparison to their Australian Caucasian counterparts. However, the absolute LLN values for DLCO and TLC did not demonstrate significant differences between some groups and the LLN values may be a better predictor for clinical decision making. Further studies are however, warranted.

Supporting information

S1 Fig. Showing aerial view of a NT Australian Indigenous community.
(PNG)

S2 Fig. Showing a NT Australian Indigenous community airstrip.
(PNG)

S3 Fig. Showing the respiratory outreach team, including respiratory specialist physician, respiratory clinical nurse consultant, respiratory trainee medical officer, respiratory and sleep technologist and the pilot helping to unload the portable DLCO equipment.
(PNG)

Acknowledgments

We thank Mr Xinlin Jing, from the Health Information Services, Royal Darwin Hospital, Darwin, Northern Territory, Australia and Mrs Joy J Mingi, Department of Public Health, Charles Darwin University, Darwin, Northern Territory, Australia for helping with lung function data collection for this study. We also sincerely thank Boehringer Ingelheim, pharmaceutical company for their generosity in voluntarily donating funding towards this study to address Indigenous Australian respiratory issues in the NT population.

Author Contributions

Conceptualization: Timothy Howarth, Helmi Ben Saad, Subash S. Heraganahally.

Data curation: Timothy Howarth, Ara J. Perez, Charmain B. Atos, Elisha White, Subash S. Heraganahally.

Formal analysis: Timothy Howarth.

Funding acquisition: Subash S. Heraganahally.

Investigation: Timothy Howarth.

Methodology: Helmi Ben Saad, Subash S. Heraganahally.

Project administration: Charmain B. Atos, Elisha White, Subash S. Heraganahally.

Resources: Ara J. Perez, Charmain B. Atos, Elisha White, Subash S. Heraganahally.

Supervision: Helmi Ben Saad, Subash S. Heraganahally.

Validation: Timothy Howarth.

Visualization: Subash S. Heraganahally.

Writing – original draft: Timothy Howarth.

Writing – review & editing: Timothy Howarth, Helmi Ben Saad, Subash S. Heraganahally.

References

1. Australian Institute of Health and Welfare 2015. The health and welfare of Australia's Aboriginal and Torres Strait Islander peoples 2015. Cat. no. IHW 147. Canberra: AIHW.
2. Kruavit A, Fox M, Pearson R, Heraganahally S. Chronic respiratory disease in the regional and remote population of the northern territory top end: A perspective from the specialist respiratory outreach service. *Aust J Rural Health*. 2017; 25:275–284. <https://doi.org/10.1111/ajr.12349> PMID: 28618108
3. Heraganahally SS, Wasgewatta SL, McNamara K, Eisemberg CC, Budd RC, Mehra S, et al. Chronic Obstructive Pulmonary Disease in Aboriginal Patients of The Northern Territory Of Australia: A Landscape Perspective. *Int J Chron Obstruct Pulmon Dis*. 2019; 14:2205–2217. <https://doi.org/10.2147/COPD.S213947> PMID: 31576117
4. Heraganahally SS, Wasgewatta SL, McNamara K, Mingi JJ, Mehra S, Eisemberg CC, et al. Chronic obstructive pulmonary disease with and without bronchiectasis in Aboriginal Australians—a comparative study. *Intern Med J*. 2019. <https://doi.org/10.1111/imj.14718>.
5. Westrupp EM, D'Esposito F, Freemantle J, Mensah FK, Nicholson JM. Health outcomes for Australian Aboriginal and Torres Strait Islander children born preterm, low birthweight or small for gestational age: A nationwide cohort study. *PloS one*. 2019; 14(2):e0212130. <https://doi.org/10.1371/journal.pone.0212130> PMID: 30785929
6. Howarth T, Brunette R, Davies T, Andrews RM, Patel BK, Tong S, et al. Antibiotic use for Australian Aboriginal children in three remote Northern Territory communities. *PloS one*. 2020; 15(4):e0231798. <https://doi.org/10.1371/journal.pone.0231798> PMID: 32302359
7. Australian Bureau of Statistics. Aboriginal and Torres Strait Islander Peoples: Smoking Trends, Australia, 1994 to 2014–15. ABS, Canberra. 4737.0. www.abs.gov.au > ausstats.

8. Carroll SJ, Dale MJ, Bailie R, Daniel M. Climatic and community sociodemographic factors associated with remote Indigenous Australian smoking rates: an ecological study of health audit data. *BMJ Open*. 2019; 9(7):e032173. <https://doi.org/10.1136/bmjopen-2019-032173> PMID: 31292187
9. Thompson BR, Stanojevic S, Abramson MJ, Beasley R, Coates A, Dent A, et al. On behalf of the ANZSRS reference ranges working party. The all-age spirometry reference ranges reflect contemporary Australasian spirometry. *Respirology*. 2011; 16: 912–917. <https://doi.org/10.1111/j.1440-1843.2011.01970.x> PMID: 21401802
10. Hall GL, Thompson BR, Stanojevic S, Abramson MJ, Beasley R, Coates A, et al. The Global Lung Initiative 2012 reference values reflect contemporary Australasian spirometry. *Respirology*. 2012; 17:1150–1. <https://doi.org/10.1111/j.1440-1843.2012.02232.x> PMID: 22849658
11. Schubert J, Kruavit A, Mehra S, Wasgeewatta S, Chang AB, Heraganahally SS. Prevalence and nature of lung function abnormalities among Indigenous Australians referred to specialist respiratory outreach clinics in the Northern Territory. *Intern Med J*. 2019; 49:217–224. <https://doi.org/10.1111/imj.14112> PMID: 30239090
12. Cooksley NA, Atkinson D, Marks GB, Toelle BG, Reeve D, Johns DP, et al. Prevalence of airflow obstruction and reduced forced vital capacity in an Aboriginal Australian population: The cross-sectional BOLD study. *Respirology*. 2015; 20:766–774. <https://doi.org/10.1111/resp.12482> PMID: 25704620
13. Hall GL, Pearson G. Reduced forced vital capacity in Aboriginal Australians: Biology or missing evidence? *Respirology*. 2015; 20:693–694. <https://doi.org/10.1111/resp.12552> PMID: 25988308
14. Heraganahally SS, Howarth T, White E, Sorger L, Binacardi E, Saad HB. Lung function parameters among “apparently healthy” Australian Aboriginal Adults: an Australian Caucasian and Global Lung Initiative (GLI-2012) various ethnic norms comparative study. *Expert Review of Respiratory Medicine*. 2020. <https://doi.org/10.1080/17476348.2021.1847649>
15. Blake TL, Chang AB, Petsky HL, Rodwell LT, Brown MG, Hill DC, et al. Spirometry reference values in Indigenous Australians: a systematic review. *Med J Aust*. 2016; 205:35–40. <https://doi.org/10.5694/mja16.00226> PMID: 27362686
16. MacIntyre N, Crapo RO, Viegi G, Johnson DC, Van der Grinten CPM, Brusasco V, et al. Standardisation of the single-breath determination of carbon monoxide uptake in the lung. *Eur Respir J*. 2005; 26: 720–735. <https://doi.org/10.1183/09031936.05.00034905> PMID: 16204605
17. Le Gouellec N, Duhamel A, Perez T, Hachulla A-L, Sobanski V, Faivre J-B, et al. Predictors of lung function test severity and outcome in systemic sclerosis-associated interstitial lung disease. *PloS One*. 2017; 12(8):e0181692. <https://doi.org/10.1371/journal.pone.0181692> PMID: 28763468
18. Kirby M, Owirangi A, Svenningsen S, Wheatley A, Coxson HO, Paterson NAM, et al. On the role of abnormal DLCO in ex-smokers without airflow limitation: symptoms, exercise capacity and hyperpolarised helium-3 MRI. *Thorax*. 2013; 68(8):752–759. <https://doi.org/10.1136/thoraxjnl-2012-203108> PMID: 23604381
19. Isaac BTJ, Thangakunam B, Cherian RA, Christopher DJ. The correlation of symptoms, pulmonary function tests and exercise testing with high-resolution computed tomography in patients with idiopathic interstitial pneumonia in a tertiary care hospital in South India. *Lung India*. 2015; 32(6):584. <https://doi.org/10.4103/0970-2113.168131> PMID: 26664164
20. Bolliger CT, Perruchoud AP. Functional evaluation of the lung resection candidate. *Eur Respir J*. 1998; 11:198–212. <https://doi.org/10.1183/09031936.98.11010198> PMID: 9543294
21. Pellegrino R, Viegi G, Brusasco V, Crapo R, Burgos F, Casaburi R, et al. Interpretative strategies for lung function tests. *Eur Respir J*. 2005; 26:948–68. <https://doi.org/10.1183/09031936.05.00035205> PMID: 16264058
22. Gibson G. Pulmonary hyperinflation a clinical overview. *Eur Respir J*. 1996; 9(12):2640–2649. <https://doi.org/10.1183/09031936.96.09122640> PMID: 8980982
23. Pellegrino R, Viegi G, Brusasco V, Crapo RO, Burgos F, Casaburi REA, et al. Interpretative strategies for lung function tests. *Eur Respir J*. 2005; 26:948–968. <https://doi.org/10.1183/09031936.05.00035205> PMID: 16264058
24. Hassel E, Stensvold D, Halvorsen T, Wisløff U, Langhammer A, Steinshamn S. Association between pulmonary function and peak oxygen uptake in elderly: the Generation 100 study. *Respiratory research*. 2015; 16(1):156. <https://doi.org/10.1186/s12931-015-0317-0> PMID: 26715058
25. Waschki B, Kirsten A, Holz O, Müller K-C, Schaper M, Sack A-L, et al. The role of sustained physical inactivity in the progression of exercise intolerance and muscle depletion in COPD. *Am J Respir Crit Care Med*. 2015; 192:295–306. <https://doi.org/10.1164/rccm.201501-0081OC> PMID: 26020495
26. Australian Institute of Health and Welfare. Australia’s mothers and babies 2017—in brief. In: AIHW, editor. Canberra 2019.

27. Ford EJ, Cade TJ, Doyle LW, Umstad MP. Pregnancy risk factors associated with birthweight of infants born to Australian Aboriginal women in an urban setting—a retrospective cohort study. *BMC Pregnancy Childbirth*. 2018; 18(1):382. <https://doi.org/10.1186/s12884-018-1946-3> PMID: 30249202
28. Heris C L, Eades SJ, Lyons L, Chamberlain C, Thomas DP. Changes in the age young Aboriginal and Torres Strait Islander people start smoking, 2002–2015. *Public health research & practice*. 2020; 30: 1–8. <https://doi.org/10.17061/phrp29121906> PMID: 32601654
29. Population—Australian Bureau of Statistics. Canberra March 2020. www.abs.gov.au › AUSSTATS.
30. Australian Bureau of Statistics. Estimates of Aboriginal and Torres Strait Islander Australians, June 2016. Canberra: ABS,; 2016.
31. Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. “ATS/ERS task force: standardisation of lung function testing” *Eur Respir J*. 2005; 26: 319–338. <https://doi.org/10.1183/09031936.05.00034805> PMID: 16055882
32. Wanger J, Clausen JL, Coates A, Pedersen OF, Brusasco V, Burgos F, et al. Standardisation of the measurement of lung volumes. “ATS/ERS task force: standardisation of lung function testing”. *Eur Respir J*. 2005; 26: 511–522. <https://doi.org/10.1183/09031936.05.00035005> PMID: 16135736
33. Gochicoa-Rangel L, Perez-Padilla R, Vazquez-García JC, Silva-Ceron M, Cid-Juarez S, Martínez-Briseno D, et al. Long-term stability of a portable carbon monoxide single-breath diffusing capacity instrument. *Respir Care*. 2017; 62:231–235. <https://doi.org/10.4187/respcare.04983> PMID: 27677305
34. Interpretation of single-breath TLC. ndd Medizintechnik AG, 03–2013, CB. AppNote-TLC-SingleBreath-V01R.docx.
35. Punjabi NM, Shade D, Wise RA. Correction of single-breath helium lung volumes in patients with airflow obstruction. *Chest*. 1998; 114:907–918. <https://doi.org/10.1378/chest.114.3.907> PMID: 9743183
36. Hankinson JL, Odencrantz JR, Fedan KB. Spirometric reference values from a sample of the general US population. *Am J Respir Crit Care Med*. 1999; 159(1):179–187. <https://doi.org/10.1164/ajrccm.159.1.9712108> PMID: 9872837
37. Prigot J. Quantification and chemical markers of tobacco-exposure. *Eur J Respir Dis*. 1987; 70:1–7. PMID: 3545884
38. Weiner DJ, Graham B, Stanojevic S. Ethnically Diverse Normative Data for Diffusing Capacity and Lung Volumes: Another Research Priority. *Ann Am Thorac Soc*. 2020; 17(1): 128. <https://doi.org/10.1513/AnnalsATS.201910-732LE> PMID: 31702941
39. Verheijden MW, Ton A, James AL, Wood M, Musk AW. Respiratory morbidity and lung function in two Aboriginal communities in Western Australia. *Respirology*. 2002; 7:247–253. <https://doi.org/10.1046/j.1440-1843.2002.00396.x> PMID: 12153691
40. Bremner PR, de Klerk NH, Ryan GF, James AL, Musk M, Murray C, et al. Respiratory symptoms and lung function in aborigines from tropical Western Australia. *Am J Respir Crit Care Med*. 1998; 158:1724–1729. <https://doi.org/10.1164/ajrccm.158.6.9702068> PMID: 9847259
41. Thompson JE, Sleight AC, Passey ME, Barnes A, Streatfield RW. Ventilatory standards for clinically well aboriginal adults. *Med J Aust*. 1992; 156:566–569. <https://doi.org/10.5694/j.1326-5377.1992.tb121420.x> PMID: 1565050
42. Musk AW, James AL, Palmer LJ, Ryan GF, Lake F, Golledge CL, et al. Respiratory infections and lung function in an Australian Aboriginal community. *Respirology*. 2008; 13:257–262. <https://doi.org/10.1111/j.1440-1843.2007.01221.x> PMID: 18339025
43. Veale A, Peat J, Salome C, Woolcock A, Thompson J. ‘Normal’ lung function in rural Australian Aborigines. *Aust N Z J Med*. 1997; 27:543–549. <https://doi.org/10.1111/j.1445-5994.1997.tb00962.x> PMID: 9404585
44. Thompson BR, Johns DP, Bailey M, Raven J, Walters EH, Abramson MJ. Prediction equations for single breath diffusing capacity (Tlco) in a middle aged caucasian population. *Thorax*. 2008; 63:889–893. <https://doi.org/10.1136/thx.2007.091959> PMID: 18390632
45. Blake TL, Chang AB, Chatfield MD, Marchant JM, McElrea MS. Global Lung Function Initiative-2012 ‘other/mixed’ spirometry reference equation provides the best overall fit for Australian Aboriginal and/or Torres Strait Islander children and young adults. *Respirology*. 2020; 25: 281–288. <https://doi.org/10.1111/resp.13649> PMID: 31339211
46. Quanjer PH, Stanojevic S, Cole TJ, Baur X, Hall GL, Culver BH, et al. Multi-ethnic reference values for spirometry for the 3–95 year age range: the global lung function 2012 equations. *Eur Respir J*. 2012; 40: 1324–1343. <https://doi.org/10.1183/09031936.00080312> PMID: 22743675
47. Stanojevic S, Graham BL, Cooper BG, Thompson BR, Carter KW, Francis RW, et al. Official ERS technical standards: Global Lung Function Initiative reference values for the carbon monoxide transfer factor for Caucasians. *Eur Respir J*. 2017; 50(3):1700010. <https://doi.org/10.1183/13993003.00010-2017> PMID: 28893868

48. White E, James A, de Klerk N, Musk A, Hall G. Selection of appropriate spirometry reference values in Aboriginal Australians. *Australian Indigenous Health Bulletin*. 2019; 19 (3).
49. Stanojevic S. Further evidence of the generalizability of the Global Lung Function Initiative reference equations for spirometry. *Respirology*. 2020; 25: 236–237. <https://doi.org/10.1111/resp.13684> PMID: 31430014
50. ndd Medical Technologies, 2017. EasyOne Pro®. <https://www.ndd.ch/en/product/easyone-pro.html> (Last access: February 20th 2021)