


RESEARCH ARTICLE

Very high prevalence of infection with the human T cell leukaemia virus type 1c in remote Australian Aboriginal communities: Results of a large cross-sectional community survey

Lloyd Einsiedel^{1*}, Hai Pham¹, Mohammad Radwanur Talukder¹, Kerry Taylor², Kim Wilson³, John Kaldor⁴, Antoine Gessain⁵, Richard Woodman⁶

1 Baker Heart and Diabetes Institute, Alice Springs, Northern Territory, Australia, **2** Poche Centre for Indigenous Health and Wellbeing, Alice Springs, Northern Territory, Australia, **3** National Serology Reference Laboratory, Melbourne, Australia, **4** Kirby Institute, University of New South Wales, Sydney, Australia, **5** Institut Pasteur, Paris, France, **6** Flinders University, Adelaide, Australia

* lloyd.einsiedel@nt.gov.au



OPEN ACCESS

Citation: Einsiedel L, Pham H, Talukder MR, Taylor K, Wilson K, Kaldor J, et al. (2021) Very high prevalence of infection with the human T cell leukaemia virus type 1c in remote Australian Aboriginal communities: Results of a large cross-sectional community survey. PLoS Negl Trop Dis 15(12): e0009915. <https://doi.org/10.1371/journal.pntd.0009915>

Editor: Masao Matsuoka, Kumamoto University: Kumamoto Daigaku, JAPAN

Received: May 3, 2021

Accepted: October 14, 2021

Published: December 8, 2021

Copyright: © 2021 Einsiedel et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: These data relate to an Australian Aboriginal population and are culturally sensitive. Although they cannot be shared publicly for ethical reasons, data will be made available to researchers who have received written approval from the Central Australian Human Research Ethics Committee (contact via cahrec@flinders.edu.au).

Abstract

Infection with the human T cell leukaemia virus type 1 (HTLV-1) subtype C is endemic among Aboriginal people in central Australia. To provide insights into the risk factors for transmission, we conducted the first large-scale, community-based prevalence study in seven remote Aboriginal communities. Residents >2 years old were invited to participate in the study between August 2014 and June 2018. HTLV-1 infection was defined as a positive western blot (WB) test or a positive HTLV-1 PCR. 720 community residents participated in the study (children <15 years, 142; adults, 578). Prevalences for children and adults were 3.5% (5/142) and 36.8% (213/578), respectively, reaching 49.3% (106/215) for those older than 45 years. A wide range of proviral loads were measured for both asymptomatic and symptomatic participants with no difference within groups according to age or gender; however, median PVL was 1.34 log₁₀ higher for symptomatic participants. The adult prevalence of HTLV-1 infection in central Australia is the highest reported worldwide. Sexual contact is likely to be the predominant mode of transmission.

Author summary

In the first large scale community study of HTLV-1 prevalence in central Australia we found an adult prevalence of 36.8%, the highest reported worldwide. Prevalence increased with age suggesting that sexual contact may be the predominant mode of transmission. Although a wide range of HTLV-1 proviral loads was found, in nearly 40% of participants this exceeded 1000 copies per 10⁵ peripheral blood leukocytes, a level that has been associated with an increased risk of chronic lung disease and death in this population. The median PVL for participants who were symptomatic with HTLV-1 associated

Funding: This project was funded by the National Health and Medical Research Council of Australia (project grant 1088517) (awarded to LE, KW, KT, and AG). The funder had no role in the study design, data collection and analysis, decision to publish, or manuscript preparation.

Competing interests: The authors have declared that no competing interests exist.

inflammatory diseases was 1.34 log₁₀ higher than that of those who were asymptomatic. These data demand a public health initiative to reduce HTLV-1 transmission among Aboriginal Australians.

Introduction

The human T cell leukaemia virus type 1 (HTLV-1) is a human retrovirus that causes a chronic infection, which preferentially involves CD4+ T cells [1]. Worldwide, HTLV-1 infects an estimated 5–10 million people, predominantly in focal geographic areas of high endemicity in southern Japan, the Caribbean basin, parts of South America and inter-tropical Africa [2]. An endemic focus is also present in central Australia where infection is with the ‘Australo-Melanesian’ HTLV-1 subtype C [2]. Transmission of HTLV-1 is via exposure to infected lymphocytes through blood contact, condomless sexual intercourse and breast-feeding [1]. Sexual transmission is thought to be the major mode of transmission in most endemic areas [3–5], including in Japan where mother-to-child transmission (MTCT) has fallen markedly in response to routine antenatal screening followed by counselling to limit or avoid breast-feeding by mothers with HTLV-1 [6].

A minority of people with HTLV-1 develop serious HTLV-1 associated complications including a rapidly progressive haematological malignancy (adult T cell leukaemia/lymphoma, ATL) [7,8] and several inflammatory disorders, the best studied of which is HTLV-1 associated myelopathy (HAM) [9,10]. Risk of these conditions is associated with a higher number of HTLV-1 infected cells in peripheral blood (HTLV-1 proviral load, PVL) [1], which was also a risk factor for pulmonary disease and gait abnormalities in a recent clinical survey in central Australia [11].

Notwithstanding recent international interest in the high prevalence of HTLV-1 infection among Aboriginal adults in central Australia [12], progress in understanding the epidemiology of HTLV-1 in this region has been slow. Infection with HTLV-1 was first reported in central Australia in 1988 [13], and each of the major recognised complications of HTLV-1 infection, including HAM and ATL, have now been reported from that region [14]. Nevertheless, actual community-based prevalence data are only available for a single small community in which 40% of adults were infected [15]. Consequently, inferences about prevalence and likely modes of transmission are derived from hospital-based [14,16] and laboratory testing series [17]. These studies have shown high seropositivity rates among Aboriginal adults and a far higher HTLV-1 seropositivity in central Australia compared to areas that are further north [17], suggesting that areas of high endemicity are concentrated in the arid interior of the country.

There has so far been no public health strategy to control HTLV-1 transmission in Australian Aboriginal communities. HTLV-1 testing is not included in routine antenatal screening [18], nor did the Australian national collaborative forum on HTLV-1 in 2018 recommend widespread HTLV-1 testing [19]. A better understanding of the patterns of occurrence of HTLV-1 at a community level is critical to guide the development of public health initiatives to reduce transmission [20]. The ethical collection of such data in a vast region that is sparsely populated by people with limited health literacy presents a considerable challenge. We now report the first large-scale, community-based, survey of HTLV-1c seroprevalence across a large area of central Australia, using a research method developed in partnership with participating communities. The study also systematically measured HTLV-1 PVL, providing an insight into relationships between HTLV-1 PVL, age and gender in this community setting.

Methods

Ethics statement

The study was developed in collaboration with Aboriginal stakeholders and primary health care providers and approved by the Central Australian Human Research Ethics Committee (HREC-14-242, HREC 15–322 and HREC 16–384). Informed written consent was obtained from all adult participants and, in the case of children, from their legal guardians.

Study setting

Central Australia comprises an area of some 1,000,000 km² in which most Aboriginal Australians reside in isolated communities under conditions of considerable socio-economic disadvantage. Most communities have populations of only several hundred people who often speak English as their second or third language. Infrastructure is limited but generally includes a small nurse-staffed, primary health care (PHC) clinic. The region is served by a single, well resourced, 186-bed hospital (Alice Springs hospital, ASH) in Alice Springs, which is the only town in the region. Seven remote communities (Fig 1) (estimated resident population (Australian census, 2016): children (<15 years), 296; adults (≥15 years), 1241) [21] participated in the study, during visits by the research team to the communities between 25 August 2014 and 30 June 2018.

For each community, the project was first discussed with community leaders, and then proposed to the community after ethics approval was granted. All Aboriginal community members >2 years old were eligible for recruitment, unless unable to give informed written consent. Consent for children was obtained from their legal guardians. Information about the




Fig 1. Map of Australia showing the study area of approximately 68,000 km² (green rectangle) within the Apatula Indigenous region (dark brown). The basemap layer was obtained from Natural Earth: <https://www.naturalearthdata.com/downloads/50m-natural-earth-2/50m-natural-earth-ii-with-shaded-relief-and-water/>.

<https://doi.org/10.1371/journal.pntd.0009915.g001>

project, and for consent, was provided in primary languages [15]. Summary data from one community have been presented previously [15].

A central survey site, which was generally the PHC clinic, was established in each community with the assistance of community members. Aboriginal researchers, fluent in local Aboriginal languages, provided transport to the survey site as needed, facilitating active recruitment and controlling participant numbers.

In order to compare PVL between asymptomatic and symptomatic participants, health records held at local clinics and at ASH were reviewed within several months of recruitment by project managers (HP, RT) who were blinded to HTLV-1 status. Records for five years prior to recruitment were reviewed, demographic information was collected and details of HTLV-1 associated conditions were recorded without regard to the date of diagnosis. Results of all chest imaging and neuroimaging from five years prior to recruitment to 30 November 2018 were reviewed at ASH (LE) blind to HTLV-1 status. The symptomatic group comprised predefined HTLV-1 related diseases: i) a haematological diagnosis of ATL, ii) radiologically defined bronchiectasis/bronchiolitis, iii) non-diabetic eye diseases diagnosed by an ophthalmologist, and documentation in the medical records of iv) myositis, v) infective dermatitis and vi) a diagnosis of 'myelopathy' or neurological symptoms associated with HTLV-1 [22,23] that were not explained by other causes. Clinical details for a subset of participants who were examined by a specialist in internal medicine (LE) have been published previously [11].

Whole blood was collected in EDTA tubes from adults and older children by venepuncture. Peripheral blood buffy coats (PBBC) and plasma were recovered and the samples stored at ASH at -80°C until testing. Collection of blood on Whatman 903 cards by finger prick using a lancet (dried blood spots, DBS) was offered to children less than approximately 10 years.

HTLV-1 serologic and molecular markers

Buffy coats and DBS were shipped to the National Serology Reference Laboratory, Melbourne. HTLV serology screening was performed on samples collected by venepuncture (662 participants) using an enzyme immune-assay (Murex HTLV I + II, DiaSorin Dartford, UK) and a particle agglutination assay (Serodia HTLV-1, Fujirebio, Tokyo, Japan). All samples reactive on either screening assay were tested by Western blot (HTLV-I/II Blot2.4, MP Diagnostics, Singapore), with the result classified as positive, indeterminate or negative according to manufacturer's criteria.

HTLV-1c PVL was assessed for all participants by real-time polymerase chain reaction (PCR) using DNA extracted from PBBC for 662 participants as previously described [15], and from DBS for 58 young children. Four 6mm discs were punched from the DBS for each participant, combined in one tube, eluted and DNA extracted using a QIAamp DNA Blood Mini Kit according to the manufacturer's instructions. For all participants, the number of copies of HTLV-1 per peripheral blood leukocytes (PBL) was then calculated and the proviral load expressed as HTLV-1 copies per 1×10^5 PBL. The lower limit of detection for PBBC samples was 6.5 copies for HTLV-1 (95% CI 5.4–8.4) and 15.6 copies for albumin (95% CI 12.9–20.0).

HTLV-1 infection was defined as a positive WB test or a positive HTLV-1 PCR test. Sufficient blood for serology was not available for the 58 young children from whom only DBS samples were collected and in these cases HTLV-1 infection was defined as a positive PCR test.

Statistical analysis

All analysis was performed using the Stata statistical software version 15.1 (StataCorp, USA). The primary outcome was HTLV-1 infection, as defined above. Categorical data were described using frequencies and percentages and continuous data using mean and standard

deviation. Differences between group means were assessed using independent t-tests and chi-squared tests of Fishers Exact test as appropriate. Logistic regression was used to determine predictors of binary outcomes with variables significant at $p < 0.20$ in univariate analysis considered for inclusion in multivariate analysis. Associations between quantities measured within the same individual were assessed using the Pearson r correlation coefficient. A 2-sided Type 1 error rate of $\alpha = 0.05$ was used for assessing statistical significance.

Results

During the study period, 720 community residents participated in the study (children < 15 , 142; adults, 578), representing 46.8% of the estimated resident population of the communities (children, 48.0%; adults 46.6%). The age and gender profile of participants closely resembled that of the overall resident population (S1 Table).

HTLV-1 infection was confirmed for 218 participants on the basis of WB or PCR positivity (WB positive/PCR positive, 209; indeterminate WB/PCR positive, 8; DBS/positive PCR, 1). There was a wide range in HTLV-1 PVL (Fig 2), but no significant difference was found between females (median PVL, 230; interquartile range (IQR) 12.0 to 1780 copies per 10^5 PBL) and males (median HTLV-1 PVL, 205; IQR 10.6 to 3630 copies per 10^5 PBL) ($p = 0.98$). There was no relationship between HTLV-1 PVL and age for females or males (Fig 2), and this was also true when symptomatic participants were excluded from the analysis (Fig 2).

Adults with HTLV-1 were more likely than participants without HTLV-1 to have conditions that were HTLV-1 associated (bronchiolitis/bronchiectasis, 27/39; possible HAM, 5/5; non-diabetic eye diseases, 4/4; infective dermatitis, 2/2; myositis, 1/1). Among 35 participants who met the definition of symptomatic HTLV-1 infection, 38 HTLV-1 associated conditions were recorded including radiologically confirmed bronchiectasis/bronchiolitis (27), possible HAM (5) (one with bronchiectasis, one with chronic bilateral corneal opacities), infective dermatitis (2) (one with bronchiectasis), non-diabetic eye diseases (4) (uveitis, 2; retinitis, 1, chronic bilateral corneal opacities, 1) and myositis (1). Evidence of neurological disease included i) gait abnormalities (three participants: a 59 year old woman with corneal opacities and urinary incontinence, PVL 45614 copies per 10^5 PBL; a 58 year old man with bronchiectasis and bladder dysfunction for whom a diagnosis of 'myelopathy' was also recorded in medical records, PVL 4160 copies per 10^5 PBL; a 57 year old woman for whom a diagnosis of 'myelopathy' was also recorded in medical records, PVL 1.88 copies per 10^5 PBL); ii) upper limb weakness, neuropathic pain, back pain and dysphagia in a 60 year old woman (PVL 6888 copies per 10^5 PBL) and iii) neuropathic pain, with back pain and bilaterally increased lower limb tone in a 44 year old man (PVL, 9100 copies per 10^5 PBL). Spinal canal stenosis was excluded by CT in four cases; no neuroimaging was done for the 59 year old woman with an abnormal gait and urinary incontinence. In no case was HTLV-1 testing done on cerebrospinal fluid (CSF).

An HTLV-1 PVL above 1000 copies per 10^5 PBL was recorded for 48 of 127 (37.8%) male participants and 31 of 91 (34.0%) female participants with HTLV-1. Notwithstanding some overlap in the PVL values of symptomatic and asymptomatic participants (Fig 2), the median HTLV-1 PVL was significantly higher for those who were symptomatic (symptomatic, 3.53, (2.65, 3.96); asymptomatic, 2.19 (0.78, 3.16) $\log_{10}$ copies per 10^5 PBL; $p < 0.001$).

The overall community HTLV-1 prevalence was 30.3% (218/720) (Table 1). Prevalences for children (under 15) and adults (15 and over) were 3.5% (5/142) and 36.8% (213/578), respectively (Table 1). None of the nineteen children aged less than 5 years had HTLV-1, and prevalence remained low for children 5–14 years (4.1%; 5/123). Thereafter, rates increased markedly, reaching 25.0% between 15–24 years and 49.3% (106/215) for adults older than 45 years (Table 1).




Fig 2. HTLV-1c proviral loads for 91 (asymptomatic, 75; symptomatic, 16) female and 128 (asymptomatic, 109; symptomatic, 19) male participants. Pearson correlation coefficients and p-values for PVL versus age are shown for symptomatic and asymptomatic subjects (symptomatic group, solid lines; asymptomatic group, dashed lines). The symptomatic group includes i) radiologically defined bronchiectasis/bronchiolitis (27) (orange circles), ii) uveitis (2), retinitis (1) or chronic bilateral corneal opacities (1) (red squares), iii) myositis (1) (green square), iv) infective dermatitis (black cross) (2) and v) neurological symptoms associated with HTLV-1 (5) (green squares). Three participants had more than one HTLV-1 associated condition: a 59 year old woman (PVL 4.65 log₁₀ copies per 10⁵ peripheral blood leukocytes, PBL) had possible HAM and eye disease (green square); a 58 year old man (PVL 3.62 log₁₀ copies per 10⁵ PBL) had possible HAM and bronchiectasis (orange circle); a 43 year old man (PVL 3.37 log₁₀ copies per 10⁵ PBL) had bronchiectasis and infective dermatitis (orange circle with black cross). Median (IQR) HTLV-1 PVL log₁₀ HTLV-1 copies per 10⁵ PBL amongst females were 2.31 (1.06–3.14) for asymptomatic females (n = 75) and 3.19 (1.73–3.89) for symptomatic females (n = 16) (p = 0.120). Median (IQR) HTLV-1 PVL log₁₀ HTLV-1 copies per 10⁵ PBL amongst males were 2.02 (0.68–3.27) for asymptomatic males (n = 109) and 3.72 (3.37, 4.28) for symptomatic males (n = 19) (p < 0.001). **Abbreviations:** BE; radiologically defined bronchiolitis/bronchiectasis; eyes, non-diabetic eye diseases; HTLV-1, human T cell leukaemia virus; IDH, infective dermatitis associated with HTLV-1; neurology, 'myelopathy' or neurological symptoms associated with HTLV-1; p, p value; PBL, peripheral blood leukocytes; r, Pearson correlation coefficient between PVL and age by gender.

<https://doi.org/10.1371/journal.pntd.0009915.g002>

As shown in Fig 3, prevalence was higher for men than women, at 38.0% (127/334) versus 23.6% (91/386); p < 0.001). Rates increased sharply for males between age groups 5–14 (4.1%) and 15–24 (37.3%) and continued to increase with age, exceeding 70% for men older than 45 years (Table 1). A more gradual increase in prevalence was apparent for women, which was most marked between age groups 15–24 (8.0%) and 25–34 years (23.5%). Thereafter, prevalence increased to 36.6% for women older than 45 years. Among women of reproductive age (15–39 years), prevalence was 20.0% (32/160).

Adult prevalence was consistently high for all seven communities, ranging from 27% to 48%. In a multivariate model that included age and gender, the odds of HTLV-1 infection increased 4% for each year of age (aOR, 1.04; 95% CI, 1.02, 1.05; p < 0.001) and was nearly three times higher for males compared to females (aOR, 2.95; 95% CI, 2.04, 4.27; p < 0.001).

Amongst participants aged 14 to 90 years old the incidence rates of HTLV-1 infection were 27.3 (22.2, 33.7) and 15.8 (10.1, 24.9) and for males and females, respectively (IRR = 1.72; 95%

Table 1. HTLV-1 prevalence by age and gender in seven remote central Australian communities, 2014–18.

Age	Female				Male				Total			
	HTLV-1 positive				HTLV-1 positive				HTLV-1 positive			
	n	(%)	OR (95% CI) ¹	p value	n	(%)	OR (95% CI) ¹	p value	n	(%)	OR (95% CI) ¹	p value
<15	3/70	4.3	ref		2/72	2.8	ref		5/142 ²	3.5	ref	
15–24	4/50	8.0	1.9 (0.4–9.1)	0.399	25/67	37.3	20.8 (4.7–92.4)	<0.001	29/117	24.8	9.0 (3.4–24.2)	<0.001
25–34	16/68	23.5	6.9 (1.9–24.8)	0.003	17/62	27.4	13.2 (2.9–60.0)	0.001	33/130	25.4	9.3 (3.5–24.7)	<0.001
35–44	19/64	29.7	9.4 (2.6–33.7)	0.001	26/52	50.0	35.0 (7.7–157.9)	<0.001	45/116	38.8	17.4 (6.6–45.7)	<0.001
45–54	26/75	34.7	11.8 (3.4–31.4)	<0.001	35/49	71.4	87.5 (18.8–406.6)	<0.001	61/124	49.2	26.5 (10.2–69.2)	<0.001
55–64	16/38	42.1	16.2 (4.3–61.0)	<0.001	13/20	65.0	65.0 (12.1–348.5)	<0.001	29/58	50.0	27.4 (9.8–76.8)	<0.001
>64	7/21	33.3	11.2 (2.6–48.6)	0.001	9/12	75.0	105.0 (15.4–715.5)	<0.001	16/33	48.5	25.8 (8.4–79.3)	<0.001
Total	91/386	23.6			127/334	38.0	2.0 (1.4–2.7)	<0.001³	218/720	30.3		

HTLV-1 infection was confirmed for 218 participants, either on the basis of WB or PCR positivity (WB positive/PCR positive, 209; indeterminate WB/PCR positive, 8; tested by dried blood spot, PCR positive, 1).

1. Unadjusted odds ratio for HTLV-1 status for male and female, and total by age groups.

2. Samples collected by dried blood spots (1/58) and PBBC (4/86).

3. Odds ratio for males relative to females

<https://doi.org/10.1371/journal.pntd.0009915.t001>

CI = 1.31, 2.27; $p < 0.001$). Amongst participants aged 20 to 24 years old the incidence rates were 10.7 (3.5, 33.2) and 2.5 (0.2, 25.1) for males and females, respectively (IRR = 4.36; 95% CI = 1.25, 15.16; $p = 0.021$). Incidence rates of HTLV-1 infection (95% CI) for asymptomatic and symptomatic men were 44.4 per 100 persons tested (35.9, 52.8) and 82.6 per 100 persons




Fig 3. HTLV-1 prevalence according to age for 386 female and 334 male community residents. HTLV-1 infection was confirmed for 218 participants, using HTLV-1 western blots (WB) and/or HTLV-1 PCR (WB positive/PCR positive, 209; indeterminate WB/PCR positive, 8; tested by dried blood spot, PCR positive, 1).

<https://doi.org/10.1371/journal.pntd.0009915.g003>

tested (45.5, 119.8), respectively. Comparable rates for asymptomatic and symptomatic women were 24.8 (19.0, 30.6) and 61.5 (31.4, 91.7), respectively. The incidence rate ratio (95% CI) of HTLV-1 infection was 1.70 (1.30, 2.24) ($p < 0.001$) for men relative to women, and was 2.10 (1.46, 3.02) ($p < 0.001$) for symptomatic relative to asymptomatic participants.

Discussion

In the first large scale, community-based, HTLV-1 prevalence study in Australia we report an adult (≥ 15 years) prevalence of 36.8%, the highest community prevalence recorded worldwide [4,24]. Micro-geographic variation in prevalence is a well-recognized feature of HTLV-1 occurrence globally [2]. For example, prior to the introduction of public health interventions that were associated with dramatic reductions in prevalence in Japan [6], rates in the south-west of that country exceeded 30% for adults aged >30 years in some villages, but were less than 10% in others nearby [24]. This was not the case in the present study in which HTLV-1 prevalence was consistently high for all participating communities across a large region of central Australia (Fig 1).

The study also systematically measured HTLV-1 PVL in study participants, extending previously published observations from these communities [11] to reveal a wide range of inter-individual HTLV-1 proviral loads, exceeding 1000 copies per 10^5 PBL for 36% of participants with HTLV-1. An HTLV-1 PVL exceeding this level has been associated with serious adverse outcomes, including bronchiectasis and death, in hospital-based studies in this population [25,26]. A strong association between high HTLV-1 PVL and chronic pulmonary disease was also found among 415 adult residents of these communities who were examined by a physician blinded to HTLV-1 status [11]. In the present study, which used medical records to identify HTLV-1 associated conditions, the median PVL for the symptomatic group was 1.34 log₁₀ higher than that of asymptomatic participants.

Higher HTLV-1 PVL among men and older participants were found in a large cohort of adults recruited from Japanese medical facilities, most of whom were tested for HTLV-1 during an illness or after a relative was found to have HTLV-1 [27]. An increasing PVL with age has therefore been suggested to contribute to the increased risk of HTLV-1 associated diseases that has been observed among older individuals [27]. In contrast, no correlation was found between HTLV-1 PVL and age for either symptomatic or asymptomatic groups in the present community-based study, nor was there any difference in median values according to gender. These findings are consistent with the results of a community survey in Gabon [28] and with studies of adults who attempted blood donation in Brazil [29], the USA [30] and Japan [31,32]. Prospective studies are necessary to more accurately define the risk of developing disease entities that is associated with a given PVL. However, ensuring that participants are correctly classified at baseline will be extremely difficult, particularly for those with bronchiolitis and early HTLV-1 associated pulmonary disease [33], and neurological disease not fulfilling HAM criteria [22,23].

As is typical of other parts of the world with endemic HTLV-1 infection [4,24,34–37], prevalence increased with age in central Australia. This relationship has previously been reported for Aboriginal people in health-care associated studies [38,39]. For example, HTLV-1 seropositivity rates among Aboriginal children (aged less than 11 years) and adults whose blood was collected for a medical audit at ASH in 1977 were 6.4% and 20.6%, respectively [38]. More recently, HTLV-1 prevalences among in-patients at ASH were 1.3% and 35.9% for children and adults, respectively [16]. In a study of patients from the north of the NT, seropositivity rates among children and adults were 0.0% and 1.65%, respectively [39].

The sharp increase in the prevalence of a chronic infection around adolescence suggests that sexual contact may be playing an important role in transmission, and this is considered to

be the explanation for the age-prevalence relationship in other HTLV-1 endemic areas [3,6,40]. The role for sexual intercourse in HTLV-1 transmission has been inferred in international studies based on associations between the prevalence of HTLV-1 infection and higher numbers of sexual partners [3,5,40–43], a longer sexual relationship with a partner at risk of HTLV-1 infection [44], the presence of genital lesions [45] and a history of sexually transmitted infections (STI), including syphilis [5,45] and gonorrhoea [5]. A previous bacterial STI was also a risk factor for HTLV-1 infection in an earlier study at ASH [16]. Genital lesions may directly increase risk of HTLV-1 infection during condomless sexual intercourse by exposing target cells to HTLV-1 infected lymphocytes in body fluids [46]. If so, the high rates of bacterial STIs [27] among young Aboriginal people in the NT might enhance the risk of HTLV-1 transmission.

Among adults, the higher prevalence that we observed in men compared to women differed from international reports, which have found a higher prevalence in women [4,24]. In the present study, we were unable to collect sensitive information relating to sexual behavior or other possible risk factors for HTLV-1 transmission, and the cause for this gender-based difference is unclear. Although condomless sexual intercourse is considered to be the major mode of transmission worldwide [3,6,40], sharing devices used for self-flagellation in a religious context was associated with HTLV-1 transmission in a recent report from the UK [47], which suggests that predisposing factors may vary in different populations. Further studies by Aboriginal researchers are needed to more clearly define the risk factors for HTLV-1 transmission in this population.

Research from Japan and other endemic countries has shown that early childhood infection is strongly associated with a longer duration of breast-feeding. This increased risk is thought to result from the continued exposure of infants to HTLV-1 infected cells in breast milk after the loss of protective maternal antibodies [35], which occurs within 12 months of age [48]. Infection rates therefore increase substantially with the duration of breast-feeding: 3.9–9.0% at 6–7 months [35,48], 20.3% after 6 months [35] and 32% if breast-feeding is continued longer than 12 months [49]. The overall childhood prevalence of 3.5% that was recorded by us is consistent with that which would result from a 20% mother-to-child transmission rate and a 20% prevalence of infection among women aged 15–39 years. Although this childhood prevalence rate is somewhat higher than that reported in other marginalised populations [4], only one positive test was recorded for a child aged less than ten years and further studies are required to determine the true rates of mother to child transmission in this region.

Strengths of this study include its community setting and the blinding of participants and researchers to HTLV-1 status until after medical records were reviewed. Nevertheless, a number of limitations must be acknowledged. First, based on ABS population data, less than 50% of the estimated resident population was recruited, raising the possibility of selection bias. We are also unable to calculate a regional HTLV-1 prevalence because only seven communities participated in the study. However, we previously reported an overall adult prevalence of 35.9% for adults admitted to ASH from across the region [16] and rates are even higher among patients from adjacent areas of the states of South Australia and Western Australia [14]. The HTLV-1 endemic area may therefore extend across a vast area of Australia that includes parts of Western Australia, South Australia and the Northern Territory. Second, HTLV-1 PCR applied to DBS samples has a lower sensitivity than it does for PBBC samples. The use of DBS samples to determine HTLV-1 status for young children may therefore underestimate HTLV-1 prevalence in this group. However, the similarly low prevalence based on PBBC specimens in the larger group of children aged 10–15 years in our survey (4.7%), and the consistency with seropositivity rates recorded for children who were in-patients at ASH in 1977 (6.4% [38]) and 2016 (1.3% [16]), gives confidence in the validity of the results. Despite the low prevalence,

acquisition of HTLV-1 infection in early life is clinically very important because of its strong association with ATL, a fatal haematological malignancy that has been reported at ASH [50]. Finally, the categorization of participants as symptomatic or asymptomatic according to diagnoses recorded in their medical records is likely to underestimate the prevalence of HTLV-1 associated diseases, and this limits our interpretation of comparisons of PVL data between these groups. For example, chest HRCT are routinely requested to further characterize abnormalities apparent on chest x-ray, the sensitivity of which is poor for the diagnosis of bronchiolitis and early bronchiectasis [51]. Consequently, chest x-rays were the only imaging modality for most participants, and 21 participants with HTLV-1 had no chest imaging done during the study period. Similarly, possible HTLV-1 associated neurology was recorded in medical records for only five of 578 (0.86%) adults in the present study compared to 15 of 415 (3.6%) adults who were identified by clinical examination [11]. In the present study, the clinical description was consistent with HAM in three cases, and an association with HTLV-1 was made more likely for two of these participants by the very high PVL and the presence of other HTLV-1 associated conditions (bronchiectasis and eye disease). Nevertheless, in no case were we able to confirm this diagnosis using medical records because these were not written with HTLV-1 associated diseases in mind, HTLV-1 testing of CSF was not done and other possible causes, such as tertiary syphilis and vitamin B12 deficiency, were not excluded. This could lead to misclassification that might affect the true difference in PVL between symptomatic and asymptomatic groups, but this is unlikely to alter our conclusion that HTLV-1 PVL did not vary with age or gender for either group. Finally, PVL were measured using DNA extracted from PBBC and the differential leukocyte count was not available for many participants in our remote study setting. HTLV-1 PVL data are therefore presented relative to the number of peripheral blood leukocytes and our data cannot be compared to that from other studies that have used peripheral blood mononuclear cells as their reference. Nevertheless, our methods have strong internal validity that has allowed clinically meaningful comparisons between symptomatic and asymptomatic participants in previous studies [11,25,26].

In the first large-scale, community-based, epidemiological study of HTLV-1 in Australia, we recorded an adult HTLV-1 prevalence of 36.8%. Based on age-specific prevalence, sexual contact appears likely to be the major mode of transmission in this region, as it is in other endemic countries [4,6]. Each of the recognised complications of HTLV-1 has been described in central Australia [14]. Infection with HTLV-1 is also associated with an increased risk of death in all endemic areas where mortality has been studied [52], and a higher HTLV-1 PVL predicts death at ASH [25,26].

Prevention is particularly important because HTLV-1 infection is lifelong and there is no treatment that directly targets the virus. Results of the present study argue for the implementation of a coordinated program to inform Aboriginal Australians of the risks posed by HTLV-1 infection and to reduce the risk of viral transmission in this population. In Japan, rising incidence rates among young adults in non-endemic, metropolitan centres have followed their migration from HTLV-1 endemic regional areas [4]. A recent report of infective dermatitis in an Aboriginal girl born in a major Australian city [53] suggests that increasing population mobility may similarly make HTLV-1 infection more relevant to Australian settings beyond the remote communities of central Australia.

Supporting information

S1 Table. Estimated resident population data not available for children aged 3–4 years (n = 18). No data for gender available for two communities (n = 102). (DOCX)

Acknowledgments

We gratefully acknowledge the contributions made to this project by Shane Schinke and Aboriginal researchers Clinton Pepperill and Una Ratara who recruited participants, PHC community clinics who generously provided space despite the heavy clinical demands made of them, and the pathology service at Alice Springs Hospital. We also thank Dr Bronwyn Silver, Central Australian Aboriginal Congress, for reviewing an earlier version of the manuscript.

Author Contributions

Conceptualization: Lloyd Einsiedel.

Data curation: Hai Pham, Mohammad Radwanur Talukder, Richard Woodman.

Formal analysis: Mohammad Radwanur Talukder, Richard Woodman.

Funding acquisition: Lloyd Einsiedel, John Kaldor, Antoine Gessain, Richard Woodman.

Investigation: Kim Wilson.

Methodology: Lloyd Einsiedel, Hai Pham, Mohammad Radwanur Talukder, John Kaldor, Antoine Gessain, Richard Woodman.

Project administration: Lloyd Einsiedel, Hai Pham, Kerry Taylor.

Resources: Lloyd Einsiedel, Hai Pham, Kerry Taylor.

Supervision: Lloyd Einsiedel, Hai Pham, Mohammad Radwanur Talukder.

Writing – original draft: Lloyd Einsiedel.

Writing – review & editing: Lloyd Einsiedel, Mohammad Radwanur Talukder, Kerry Taylor, Kim Wilson, John Kaldor, Antoine Gessain, Richard Woodman.

References

1. Verdonck K, Gonzalez E, Van Dooren S, Vandamme A, Vanham G, Gotuzzo E. Human T-lymphotropic virus 1: recent knowledge about an ancient infection. *Lancet Infect Dis*. 2007; 7:266–281. [https://doi.org/10.1016/S1473-3099\(07\)70081-6](https://doi.org/10.1016/S1473-3099(07)70081-6) PMID: 17376384
2. Gessain A, Cassar O. Epidemiological aspects and world distribution of HTLV-1 infection. *Front Microbiol*. 2012; 3:388. <https://doi.org/10.3389/fmicb.2012.00388> PMID: 23162541
3. Murphy E L WR, Hanchard B, Cranston B, Figueroa J P, Gibbs W N, Murphy J et al. A case-control study of risk factors for seropositivity to human T-lymphotropic virus type I (HTLV-I) In Jamaica. *Int. J. Epidemiol*. 1996; 25(5):1083–1089. <https://doi.org/10.1093/ije/25.5.1083> PMID: 8921498
4. Plancoulaine S, Buigues RP, Murphy EL, van Beveren M, Pouliquen JF, Joubert M et al. Demographic and familial characteristics of HTLV-I infection among an isolated, highly endemic population of African origin in French Guiana. *Int J Cancer*. 1998 May 4; 76(3):331–6. [https://doi.org/10.1002/\(sici\)1097-0215\(19980504\)76:3<331::aid-ijc8>3.0.co;2-w](https://doi.org/10.1002/(sici)1097-0215(19980504)76:3<331::aid-ijc8>3.0.co;2-w) PMID: 9579568
5. Nakashima K, Kahiwa S., Kajiyama W., Hirata M., Hayashi J., Noguchi A et al. Sexual transmission of Human T-lymphotropic virus Type I among female prostitutes and among patients with sexually transmitted diseases in Fukuoka, Kyushu, Japan. *Am J Epidemiol*. 1995; 141:305–311. <https://doi.org/10.1093/aje/141.4.305> PMID: 7840108
6. Satake M, Iwanaga M, Sagara Y, Watanabe T, Okuma K, Isao Hamaguchi I. Incidence of human T-lymphotropic virus 1 infection in adolescent and adult blood donors in Japan: a nationwide retrospective cohort analysis. *Lancet Infect Dis*. 2016; 16:1246–1254. [https://doi.org/10.1016/S1473-3099\(16\)30252-3](https://doi.org/10.1016/S1473-3099(16)30252-3) PMID: 27567105
7. Uchiyama T, Yodoi J, Sagawa K, Takatsuki K, Uchino H. Adult T-cell leukemia: clinical and hematologic features of 16 cases. *Blood*. 1977 Sep; 50(3):481–92. PMID: 301762
8. Poesz BJ, Ruscetti FW, Gazdar AF, Bunn PA, Minna JD, Gallo RC. Detection and isolation of type C retrovirus particles from fresh and cultured lymphocytes of a patient with cutaneous T-cell lymphoma.

- Proc Natl Acad Sci U S A. 1980 Dec; 77(12):7415–9. <https://doi.org/10.1073/pnas.77.12.7415> PMID: 6261256
9. Gessain A, Barin F, Vernant JC, Gout O, Maurs L, Calender A, de Thé G. Antibodies to human T-lymphotropic virus type-I in patients with tropical spastic paraparesis. *Lancet*. 1985 Aug 24; 2(8452):407–10. [https://doi.org/10.1016/s0140-6736\(85\)92734-5](https://doi.org/10.1016/s0140-6736(85)92734-5) PMID: 2863442
 10. Osame M, Usuku K, Izumo S, Ijichi N, Amitani H, Igata A, Matsumoto M, Tara M. HTLV-I associated myelopathy, a new clinical entity. *Lancet*. 1986 May 3; 1(8488):1031–2. [https://doi.org/10.1016/s0140-6736\(86\)91298-5](https://doi.org/10.1016/s0140-6736(86)91298-5) PMID: 2871307
 11. Einsiedel L, Pham H, Talukder MRR, Liddle J, Taylor K, Wilson K, Jersmann H, Gessain A, Woodman R, Kaldor J. Pulmonary disease is associated with human T-cell leukaemia virus type 1c infection: a cross-sectional survey in remote Aboriginal Australian communities. *Clin Infect Dis*. 2020 Sep 16: ciae1401. <https://doi.org/10.1093/cid/ciae1401> PMID: 32936911
 12. Martin F, Tagaya Y, Gallo R. Time to eradicate HTLV-1: an open letter to WHO. *The Lancet*. 2018 May 12; 391(10133):1893–4. [https://doi.org/10.1016/S0140-6736\(18\)30974-7](https://doi.org/10.1016/S0140-6736(18)30974-7) PMID: 29781438
 13. May J, Stent G, Schnagl R. Antibody to human T-cell lymphotropic virus type 1 in Australian Aborigines. *Med J Aust*. 1988; 149:104.
 14. Einsiedel L, Spelman T, Goeman E, Cassar O, Arundell M, Gessain A. Clinical Associations of Human T-Lymphotropic Virus Type 1 Infection in an Indigenous Australian Population. *PLoS Negl Trop Dis* 2014 8(1):e2643 <https://doi.org/10.1371/journal.pntd.0002643> PMID: 24454973
 15. Einsiedel L, Pham H, Woodman R, Pepperill C, Taylor K. The prevalence and clinical associations of HTLV-1 infection in a remote Indigenous community. *Med J Aust*. 2016; 205(7):305–309. <https://doi.org/10.5694/mja16.00285> PMID: 27681971
 16. Einsiedel L, Woodman R, Flynn M, Wilson K, Cassar O, Gessain A. Human T-Lymphotropic Virus type 1 infection in an Indigenous Australian population: epidemiological insights from a hospital-based cohort study. *BMC Public Health*. 2016; 16:787. <https://doi.org/10.1186/s12889-016-3366-5> PMID: 27526923
 17. Grivas R, Freeman K, Human R. B.T-lymphotropic virus-1 serology in the Northern Territory: 2008–2011. *Pathology*. 2014; 46(7):644–648. <https://doi.org/10.1097/PAT.000000000000164> PMID: 25393257
 18. Council CAaNH. *Minymaku Kutju Tjukurpa—Women's Business Manual 5th Edition* Alice Springs: Central Australian Rural Practitioner's Association; 2014.
 19. Department of Health. Paper presented at: HTLV-1 collaborative forum 2018; Alice Springs.
 20. Gordon C HS. Current knowledge about HTLV-1 in Central Australia: Proceedings from the first workshop on HTLV-1 in Central Australia. *Northern Territory Disease Control Bulletin*. 2012; 19(March):14.
 21. Australian Bureau of Statistics. *Themes: QuickStats*, viewed December 2019, <https://www.abs.gov.au/websitedbs/D3310114.nsf/Home/2016%20QuickStats>
 22. Araujo AQ, Silva MT. The HTLV-1 neurological complex. *The Lancet Neurology*. 2006 Dec 1; 5(12):1068–76. [https://doi.org/10.1016/S1474-4422\(06\)70628-7](https://doi.org/10.1016/S1474-4422(06)70628-7) PMID: 17110288
 23. Araujo AQC, Wedemann D. HTLV-1 Associated Neurological Complex. What is Hidden below the Water? *AIDS Rev*. 2019; 21(4):211–217. <https://doi.org/10.24875/AIDSRev.19000108> PMID: 31834313
 24. Tajima K, Kamura S, Ito S-I, Nagatomo M, Kinoshita K-I, Ikeda S. Epidemiological features of HTLV-1 carriers and incidence of ATL in an ATL-endemic island: a report of the community-based co-operative study in Tsushima, Japan. *Int J Cancer*. 1987; 40:741–746. <https://doi.org/10.1002/ijc.2910400605> PMID: 2891624
 25. Einsiedel L, Pham H, Wilson K, et al. Human T-Lymphotropic Virus type 1c subtype proviral loads, chronic lung disease and survival in a prospective cohort of Indigenous Australians. *PLoS Negl Trop Dis*. 2018; 12(3):e0006281. <https://doi.org/10.1371/journal.pntd.0006281> PMID: 29529032
 26. Einsiedel L, Pham H, Au V, et al. Predictors of non-cystic fibrosis bronchiectasis in Indigenous adult residents of central Australia: results of a case-control study. *ERJ Open Res*. 2019;5. <https://doi.org/10.1183/23120541.00001-2019> PMID: 31911928
 27. Iwanaga M, Watanabe T, Utsunomiya A, Okayama A, Uchimaru K, Koh KR, Ogata M, Kikuchi H, Sagara Y, Uozumi K, Mochizuki M. Human T-cell leukemia virus type I (HTLV-1) proviral load and disease progression in asymptomatic HTLV-1 carriers: a nationwide prospective study in Japan. *Blood*. 2010 Aug 26; 116(8):1211–9. <https://doi.org/10.1182/blood-2009-12-257410> PMID: 20448111
 28. Caron M, Besson G, Padilla C, Makuwa M, Nkoghé D, Leroy E, Kazanji M. Revisiting human T-cell lymphotropic virus types 1 and 2 infections among rural population in Gabon, central Africa thirty years after the first analysis. *PLoS neglected tropical diseases*. 2018 Oct 25; 12(10):e0006833. <https://doi.org/10.1371/journal.pntd.0006833> PMID: 30359373

29. Furtado MD, Andrade RG, Romanelli LC, Ribeiro MA, Ribas JG, Torres EB, Barbosa-Stancioli EF, Proietti AB, Martins ML. Monitoring the HTLV-1 proviral load in the peripheral blood of asymptomatic carriers and patients with HTLV-associated myelopathy/tropical spastic paraparesis from a Brazilian cohort: ROC curve analysis to establish the threshold for risk disease. *Journal of medical virology*. 2012 Apr; 84(4):664–71. <https://doi.org/10.1002/jmv.23227> PMID: 22337307
30. Murphy EL, Lee TH, Chafets D, Nass CC, Wang B, Loughlin K, Smith D. Higher human T lymphotropic virus (HTLV) provirus load is associated with HTLV-I versus HTLV-II, with HTLV-II subtype A versus B, and with male sex and a history of blood transfusion. *Journal of Infectious Diseases*. 2004 Aug 1; 190(3):504–10.
31. Nagai M, Usuku K, Matsumoto W, Kodama D, Takenouchi N, Moritoyo T, Hashiguchi S, Ichinose M, Bangham CR, Izumo S, Osame M. Analysis of HTLV-I proviral load in 202 HAM/TSP patients and 243 asymptomatic HTLV-I carriers: high proviral load strongly predisposes to HAM/TSP. *J Neurovirol*. 1998 Dec; 4(6):586–93. <https://doi.org/10.3109/13550289809114225> PMID: 10065900
32. Etoh K, Yamaguchi K, Tokudome S, Watanabe T, Okayama A, Stuver S, Mueller N, Takatsuki K, Mat-suoka M. Rapid quantification of HTLV-I provirus load: detection of monoclonal proliferation of HTLV-I-infected cells among blood donors. *Int J Cancer*. 1999 Jun 11; 81(6):859–64. [https://doi.org/10.1002/\(sici\)1097-0215\(19990611\)81:6<859::aid-ijc4>3.0.co;2-k](https://doi.org/10.1002/(sici)1097-0215(19990611)81:6<859::aid-ijc4>3.0.co;2-k) PMID: 10362130
33. Einsiedel L, Chiong F, Jersmann H, Taylor GP. Human T-cell leukaemia virus type 1 associated pulmonary disease: clinical and pathological features of an under-recognised complication of HTLV-1 infection. *Retrovirology*. 2021 Jan 6; 18(1):1. Erratum in: *Retrovirology*. 2021 Feb 22; 18(1):5. <https://doi.org/10.1186/s12977-020-00543-z> PMID: 33407607
34. Larsen O, Andersson S, da Silva Z, Hedegaard K, Sandström A, Naclér A, Dias F, Melbye M, Aaby P. Prevalences of HTLV-1 infection and associated risk determinants in an urban population in Guinea-Bissau, West Africa. *Journal of acquired immune deficiency syndromes (1999)*. 2000 Oct 1; 25(2):157–63.
35. Hino S. Establishment of the milk-borne transmission as a key factor for the peculiar endemicity of human T-lymphotropic virus type 1 (HTLV-1): the ATL Prevention Program Nagasaki. *Proceedings of the Japan Academy, Series B*. 2011 May 11; 87(4):152–66. <https://doi.org/10.2183/pjab.87.152> PMID: 21558754
36. Alarcón JO, Friedman HB, Montano SM, Zunt JR, Holmes KK, Quinnan GV Jr. High endemicity of human T-cell lymphotropic virus type 1 among pregnant women in Peru. *Journal of acquired immune deficiency syndromes (1999)*. 2006 Aug 15; 42(5):604. <https://doi.org/10.1097/01.qai.0000221680.52563.d5> PMID: 16773029
37. Delaporte E, Dupont A, Peeters M, Josse R, Merlin M, Schrijvers D, Hamono B, Bedjabaga L, Cherin-gou H, Boyer F, Brun-Vézinet F. Epidemiology of HTLV-I in Gabon (western equatorial Africa). *International journal of cancer*. 1988 Nov 15; 42(5):687–9. <https://doi.org/10.1002/ijc.2910420509> PMID: 3182104
38. May JT, Stent G, Bishop F, D. S. Prevalence of antibody to human T-lymphotropic virus type-1 (HTLV-1) in Australian aborigines, and detection in Indonesian sera. *Acta Virol*. 1990; 34(1):80–84. PMID: 1975728
39. Bastian I. *HTLV-1 studies in the Northern Territory*. Darwin: Menzies School of Health Research, University of Sydney; 1996.
40. Nunes D, Boa-Sorte N, Grassi MF, Taylor GP, Teixeira MG, Barreto ML, Dourado I, Galvão-Castro B. HTLV-1 is predominantly sexually transmitted in Salvador, the city with the highest HTLV-1 prevalence in Brazil. *PloS one*. 2017 Feb 3; 12(2):e0171303. <https://doi.org/10.1371/journal.pone.0171303> PMID: 28158226
41. Figueroa JP, Morris J, Brathwaite A, Ward E, Peruga A, Hayes R, Vermund SH, Blattner W. Risk factors for HTLV-I among heterosexual STD clinic attenders. *Journal of acquired immune deficiency syndromes and human retrovirology: official publication of the International Retrovirology Association*. 1995 May 1; 9(1):81–8. PMID: 7712238
42. Schreiber GB, Murphy EL, Horton JA, Wright DJ, Garfein R, Chien HC, Nass CC. Risk factors for human T-cell lymphotropic virus types I and II (HTLV-I and-II) in blood donors: the Retrovirus Epidemiology Donor Study. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 1997 Mar 1; 14(3):263–71. <https://doi.org/10.1097/00042560-199703010-00011> PMID: 9117460
43. Sanchez-Palacios C, Gotuzzo E, Vandamme AM, Maldonado Y. Seroprevalence and risk factors for human T-cell lymphotropic virus (HTLV 1) infection among ethnically and geographically diverse Peruvian women. *International journal of infectious diseases*. 2003 Jun 1; 7(2):132–7. [https://doi.org/10.1016/s1201-9712\(03\)90009-9](https://doi.org/10.1016/s1201-9712(03)90009-9) PMID: 12839715
44. Kaplan JE, Khabbaz RF, Murphy EL, Hermansen S, Roberts C, Lal R, Heneine W, Wright D, Matijas L, Thomson R, Rudolph D. Male-to-female transmission of human T-cell lymphotropic virus types I and II:

- association with viral load. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 1996 Jun 1; 12(2):193–201. <https://doi.org/10.1097/00042560-199606010-00014> PMID: 8680892
45. Murphy E FJ, Gibbs WN, et al. Sexual transmission of human T-lymphotropic virus type I (HTLV-I). *Ann. Intern. Med.* 1989; 111:555–560. <https://doi.org/10.7326/0003-4819-111-7-555> PMID: 2789009
 46. Paiva A, Casseb J. Sexual transmission of human T-cell lymphotropic virus type 1. *Revista da Sociedade Brasileira de Medicina Tropical*. 2014 Jun; 47(3):265–74. <https://doi.org/10.1590/0037-8682-0232-2013> PMID: 25075475
 47. Tang AR, Taylor GP, Dhasmana D. Self-Flagellation as Possible Route of Human T-Cell Lymphotropic Virus Type-1 Transmission. *Emerg Infect Dis.* 2019 Apr; 25(4):811–813. <https://doi.org/10.3201/eid2504.180984> PMID: 30882326
 48. Rosadas C, Taylor GP. Mother-to-child HTLV-1 transmission: unmet research needs. *Frontiers in microbiology*. 2019 May 8; 10:999. <https://doi.org/10.3389/fmicb.2019.00999> PMID: 31134031
 49. Wiktor SZ, Pate EJ, Rosenberg PS, Barnett M, Palmer P, Medeiros D, Maloney EM, Blattner WA. Mother-to-child transmission of human T-cell lymphotropic virus type I associated with prolonged breast-feeding. *Journal of human virology*. 1997 Nov 1; 1(1):37–44. PMID: 10195229
 50. Einsiedel L, Cassar O, Bardy P, Kearney D, Gessain A. Variant Human T-cell Lymphotropic Virus Type 1c and Adult T-cell Leukemia, Australia. *Emerg Infect Dis.* 2013; 19(10):1639–1641. <https://doi.org/10.3201/eid1910.130105> PMID: 24047544
 51. Einsiedel L, Fernandes L, Spelman T, Steinfort D, Gotuzzo E. Bronchiectasis is associated with human T-lymphotropic virus 1 infection in an Indigenous Australian population. *Clin Infect Dis.* 2012 Jan 1; 54(1):43–50. <https://doi.org/10.1093/cid/cir766> Epub 2011 Nov 17. PMID: 22095566.
 52. Schierhout G, McGregor S, Gessain A, Einsiedel L, Martinello M, Kaldor J. Association between HTLV-1 infection and adverse health outcomes: a systematic review and meta-analysis of epidemiological studies. *Lancet Infect Dis.* 2019. [https://doi.org/10.1016/S1473-3099\(19\)30402-5](https://doi.org/10.1016/S1473-3099(19)30402-5) PMID: 31648940
 53. Menz F, Menz J, Wilson K, Turpin J, Bangham C, Einsiedel L. Infective dermatitis associated with human T-lymphotropic virus type 1 infection in Adelaide, South Australia. *Australas J Dermatol.* 2018 May; 59(2):151–153. <https://doi.org/10.1111/ajd.12704> Epub 2017 Aug 22. PMID: 28832919.