

RESEARCH ARTICLE

The efficacy of sampling strategies for estimating scabies prevalence

Nefel Tellioglu¹, Rebecca H. Chisholm^{2,3}, Jodie McVernon^{3,4}, Nicholas Geard^{1,5}, Patricia Therese Campbell^{3,4*}

1 School of Computing and Information Systems, The University of Melbourne, Melbourne, Australia, **2** Department of Mathematics and Statistics, La Trobe University, Bundoora, Australia, **3** Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, Australia, **4** Department of Infectious Diseases, University of Melbourne, at the Peter Doherty Institute for Infection and Immunity, Melbourne, Australia, **5** Department of Infectious Diseases, The University of Melbourne, Melbourne, Australia

* patricia.campbell@unimelb.edu.au



Abstract

Background

Estimating community level scabies prevalence is crucial for targeting interventions to areas of greatest need. The World Health Organisation recommends sampling at the unit of households or schools, but there is presently no standardised approach to scabies prevalence assessment. Consequently, a wide range of sampling sizes and methods have been used. As both prevalence and drivers of transmission vary across populations, there is a need to understand how sampling strategies for estimating scabies prevalence interact with local epidemiology to affect the accuracy of prevalence estimates.

Methods

We used a simulation-based approach to compare the efficacy of different scabies sampling strategies. First, we generated synthetic populations broadly representative of remote Australian Indigenous communities and assigned a scabies status to individuals to achieve a specified prevalence using different assumptions about scabies epidemiology. Second, we calculated an observed prevalence for different sampling methods and sizes.

Results

The distribution of prevalence in subpopulation groups can vary substantially when the underlying scabies assignment method changes. Across all of the scabies assignment methods combined, the simple random sampling method produces the narrowest 95% confidence interval for all sample sizes. The household sampling method introduces higher variance compared to simple random sampling when the assignment of scabies includes a household-specific component. The school sampling method overestimates community prevalence when the assignment of scabies includes an age-specific component.

OPEN ACCESS

Citation: Tellioglu N, Chisholm RH, McVernon J, Geard N, Campbell PT (2022) The efficacy of sampling strategies for estimating scabies prevalence. *PLoS Negl Trop Dis* 16(6): e0010456. <https://doi.org/10.1371/journal.pntd.0010456>

Editor: Roger Salamon, INSERM U897, FRANCE

Received: December 15, 2021

Accepted: April 30, 2022

Published: June 9, 2022

Copyright: © 2022 Tellioglu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All data are available online. Household-age structure data is available at: <https://doi.org/10.7717/peerj.3958/supp-3> Other data are available at https://github.com/nefeltellioglu/sampling_strategy.

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Discussion

Our results indicate that there are interactions between transmission assumptions and surveillance strategies, emphasizing the need for understanding scabies transmission dynamics. We suggest using the simple random sampling method for estimating scabies prevalence. Our approach can be adapted to various populations and diseases.

Author summary

Scabies is a parasitic infestation that is commonly observed in disadvantaged populations. A wide range of sampling sizes and methods have been used to estimate scabies prevalence. With differing key drivers of transmission and varying prevalence across populations, it can be challenging to determine an effective sampling strategy. In this study, we propose a simulation approach to compare the efficacy of different sampling methods and sizes. First, we generate synthetic populations and then assign a scabies status to individuals to achieve a specified prevalence using different assumptions about scabies epidemiology. Second, we calculate an observed prevalence for different sampling methods and sizes. Our results indicate that there are interactions between transmission assumptions and surveillance strategies. We suggest using the simple random sampling method for estimating prevalence as it produces the narrowest 95% confidence interval for all sampling sizes. We propose guidelines for determining a sample size to achieve a desired level of precision in 95 out of 100 samples, given estimates of the population size and a priori estimates of true prevalence. Our approach can be adapted to various populations, informing an appropriate sampling strategy for estimating scabies prevalence with confidence.

Introduction

Scabies is a parasitic infestation caused by the mite *Sarcoptes scabiei* [1] and is one of the highest-burden Neglected Tropical Diseases (NTDs) [2]. In 2016, it was estimated that scabies affects 455 million people annually and causes 3.8 million disability-adjusted life years (DALYs) [2]. The prevalence of scabies is highest in underprivileged tropical settings including Indigenous communities of Australia and Pacific Island communities [3–5]. These settings are thought to be affected due to factors such as overcrowding [6], hot weather and humidity [7, 8]. Scabies prevalence can reach up to 35% in remote Indigenous communities [9, 10] and 71% in Pacific Island communities [11]. In these humid low-income settings, the scratching due to scabies can lead to secondary skin infections by Group A *Streptococcus* and their sequelae, which scales up the burden of scabies [8, 12]. Interventions for controlling scabies can reduce the burden of not only scabies but also the secondary skin infections [8]. Such interventions can be costly to implement [4] and it is desirable to focus such efforts on high prevalence settings.

A recent report of the World Health Organization (WHO) Informal Consultation on a Framework for Scabies Control suggests local scabies prevalence of ten percent or more as a threshold for implementing mass drug administration (MDA) of ivermectin/permethrin [13]. For this intervention to be applied appropriately, an accurate estimation of true prevalence in communities is crucial. For prevalence estimation, WHO recommends community-based household sampling methods including people from all ages as the most appropriate

strategy. They also suggest that school-based sampling might be an alternative, however, note further research is needed to determine how school scabies prevalence is related to community prevalence [13]. Even though the WHO recommends sampling strategies for estimating scabies prevalence, they underline the need for evaluation of the efficacy of such strategies. The design of studies that are both efficient and unbiased can be challenging due to following key issues.

First, the extent to which standard sample size calculation formulas are applicable to scabies is uncertain, as substantial heterogeneity in prevalence is observed between households and age groups. Prevalence studies have shown that household contacts play a crucial role in scabies transmission [14, 15]. It is estimated that it takes on average 20 minutes of close contact for scabies transmission [14] which highlights household contacts as an important factor in scabies transmission [14, 16, 17]. Dagne et al. [15] found that the probability of being infested by scabies was almost five times higher among participants with at least one household member having an itchy lesion than participants without family members with such a lesion, underlying the crucial role of the household in transmission. Other prevalence studies conducted in different settings have found scabies prevalence to be age-dependent, with children experiencing prevalence around two to three times that of adults [11, 18–23].

Second, it is hard to compare effectiveness of different sampling strategies across populations. Prevalence estimation studies have used different sampling strategies due to the varying reasons such estimates were required (for example estimating the prevalence in schoolchildren or applying mass drug administration (MDA)) [15, 18, 24–28]. For example, to estimate the level of treatment uptake in households with clinically diagnosed scabies cases, La et al. [27] screened households based on previous enrolment into a related study and found that 23% of the screened population had scabies before the intervention. To estimate scabies prevalence in a welfare home in Malaysia, Zayyid et al. [28] screened a random selection of 120 out of 160 children and found 31% of children had scabies. Moreover, highly variable scabies prevalence has been observed in survey studies conducted in Australian Indigenous communities (from 5% to 35%) [11], and other Pacific Island communities (from 5% to 71%) [11]. With no standardization in methods for scabies prevalence estimation [4], it is difficult to make valid comparisons across settings.

In order to evaluate sampling strategies for infectious disease prevalence, there are published simulation-based approaches [29–35]. Such simulation-based approaches allow us to introduce disease and population-specific characteristics and conduct *in silico* experiments on the effectiveness of sampling strategies. For example, Giardina et al. [29] used a dynamic simulation to compare efficacy of sampling strategies for monitoring morbidity targets for soil-transmitted helminths in districts consisting of villages. They found that sampling school-aged children from ten instead of five villages would increase the sampling effectiveness by 20%. Schmidt et al. [30] found that clustering among individuals and infection duration were major factors contributing to the effectiveness of sampling strategies to measure the prevalence of recurrent infections.

With the uncertainty around age-dependent prevalence and level of household transmission, it remains to be determined whether common sampling approaches introduce a bias in estimating the true prevalence of scabies. In this study, we evaluated the efficacy of different sampling strategies to estimate scabies prevalence using a simulation-based approach. We demonstrate our approach in the context of remote Australian Indigenous communities. Our approach allows comparison of the performance of sampling strategies in a simulated population which has similar age and household size distributions to Australian Indigenous communities.

Methods

In order to compare and evaluate different sampling strategies, we simulated a population with a known true prevalence of scabies, and also simulated the sampling strategy used to estimate the prevalence of scabies in that population. By comparing the estimated prevalence to the true prevalence, we evaluated the efficacy of a given sampling strategy and compare the performance of different strategies. Our approach consists of three stages. First, we generated synthetic populations with characteristics similar to those of remote Australian Indigenous communities. Second, given uncertainty around the relative importance of age- and household-specific factors, we examined five different rules for attributing disease status in the population. We assigned a positive or negative scabies status to all individuals in the population to achieve a specified prevalence. Third, we sampled a percentage of this synthetic population using a pre-defined sampling strategy and sample size and record the sample prevalence. Finally, we compared the specified prevalence (the input prevalence) with the sample prevalence (the output prevalence).

Generating synthetic populations

We generated populations ranging in size from approximately 500 to 4000 individuals representing the population size of medium to large remote Australian Indigenous communities [36]. Within each population, we assigned individuals into households and age-classes. Age-classes consist of *adult* (16 years and over), *school* (5–15 years), and *pre-school* (0–4 years) at random such that the household size distribution and age distribution of the population reflected Australian Bureau of Statistics (ABS) 2016 census data [37] and survey data of Vino et al. [38].

To generate a population of size N , we repeatedly sampled household sizes from the household size distribution of Indigenous communities in ABS data until the population contained approximately N people (with a tolerance of 5%). In ABS data, the distribution of households having a size of six or larger is aggregated. After sampling household sizes ranging [1, . . . , 5, 6+], we used survey data of Vino et al. [38] to disaggregate data for households with a size of six or larger. Household size distribution of all of the simulated populations (blue) and ABS data (red) are represented in Fig 1.

For each household, given a household size, we sampled age-classes (*adult*, *school*, *pre-school*) for household members based on age-class distribution of household data from [38]. We repeated the sampling of age-classes for each household until it contained at least one adult. In the survey data, there was no age-class distribution for households of size 15, 18, 19, 20, and 22. For these households, we used the age-class distribution of the closest household size. We then compared the age-class structure of all of the simulated populations with independent age structure data from the Aboriginal and Torres Strait Islander Health Performance Framework 2017 [39] (Fig 2).

Assigning scabies status

For each of the generated populations, we assigned a positive or negative scabies status to individuals to achieve an input prevalence percentage, ranging from 5% to 40%. The *assignment method* used to set the scabies status of individuals was chosen from one of the following:

1. **Random:** Individuals were assigned a positive scabies status uniformly at random.
2. **Household-specific (high):** Households were selected uniformly at random, and all individuals in the selected households were assigned a positive scabies status.




Fig 1. Household size distribution. Household size distribution (median and 2.5–97.5 quantiles) of simulated data (blue) and Indigenous communities household size distribution taken from Australian Bureau of Statistics 2016 census data (red) [37] are presented.

<https://doi.org/10.1371/journal.pntd.0010456.g001>




Fig 2. Age distribution. Age distribution (median and 2.5–97.5 quantiles) of simulated data (blue) and age distribution taken from Aboriginal and Torres Strait Islander Health Performance Framework 2017 (red) [39] are presented.

<https://doi.org/10.1371/journal.pntd.0010456.g002>

3. **Household-specific (mild):** Households were selected uniformly at random, and half of the individuals in the selected households were assigned a positive scabies status, uniformly at random.
4. **Age-specific:** Individuals were assigned a positive scabies status uniformly at random; however, children were three times more likely than adults to be assigned a positive scabies status, based on scabies prevalence surveys conducted in NT, Fiji, and Ethiopia [19–21].
5. **Age-and-household-specific:** Households were selected uniformly at random, and a positive scabies status was assigned to half of the individuals in selected households, with children three times more likely than adults to be assigned a positive scabies status.

Simulating sampling strategies

We simulated three different *sampling methods* in each of the generated populations: random, household or school sampling with *sample sizes* between 5% and 90% of the population, assuming all individuals were available for sampling. In this study, we refer to a combination of a sampling method and a sampling size as a *sampling strategy*. The simple random sampling method involved sampling individuals uniformly at random. The household sampling method involved selecting households uniformly at random and sampling all members. The school sampling method involved sampling individuals uniformly at random from the school age group only.

Study design

A population is generated with a size sampled uniformly at random in the range of 500 to 4000. For each generated population with age and household structure, we assign scabies with one of the five assignment methods followed by sampling with the chosen strategy (Table 1). For school sampling, where it is not possible to sample the chosen sample size due to the size of the schoolchildren population, we stop sampling when all schoolchildren have been selected. This process is repeated 500 times. A simplified pseudo code is provided in Fig 3.

In order to compare different sampling methods, we first calculated scabies prevalence distribution in age and household groups given a scabies assignment method and input prevalence. Then, as an exemplar, we compared the output prevalence in the samples using different sampling strategies when the input prevalence was between 20–30%. Finally, we calculated the sample size required to achieve a target precision for each sampling method under different population size, prevalence, and assignment method scenarios.

Table 1. Model Parameters.

Symbol	Description	Values	Reference
M	Number of simulations	500	-
N	Population size	\sim Uniform(500, 4000)	[36]
-	Specified prevalence percentage (%)	5, 10, 15, 20, 25, 30, 35, 40	[9, 11, 20]
-	Scabies assignment method	“random”, “high household-specific”, “mild household-specific”, “age-specific”, “age-and-household-specific”	[11, 14, 15, 19]
-	Sampling method	“random”, “household”, “school”	[13, 15, 18]
-	Sampling percentage (%)	5, 10, 15, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90	-

<https://doi.org/10.1371/journal.pntd.0010456.t001>

```

1 for  $i \in \{1, 2, \dots, M\}$  do
2   Generate a population size,  $N$ ;
3   Generate a population dataset, with size of  $N$  and a size tolerance of 5% by
   assigning household and age groups;
4   for each assignment method do
5     for each specified prevalence percentage do
6       Assign scabies to selected individuals based on assignment method
       and specified prevalence percentage;
7       for each sampling method do
8         for each sampling percentage do
9           Sample the population using the sampling method and
           sampling percentage;
10        end
11       end
12     end
13   end
14 end

```

Fig 3. Pseudo code of our algorithm.

<https://doi.org/10.1371/journal.pntd.0010456.g003>

Results

We observed that the distribution of prevalence in population groups can vary substantially when the underlying scabies assignment method changes. For example, age-specific scabies assignment increases the prevalence among children as well as prevalence in larger households, due to the higher number of children in larger households (Fig 4A and 4B). In addition, household specific assignment approaches introduce higher variance in prevalence among households (Fig 5).




Fig 4. (A) Distribution of scabies prevalence in age groups for different scabies assignment methods, (B) Distribution of scabies prevalence across household size groups for different methods of scabies status assignment. The results (median and 2.5% to 97.5% quantiles) are plotted for an exemplar input prevalence percentage between 20–30% across all population sizes. In the legend, “HH” refers to household.

<https://doi.org/10.1371/journal.pntd.0010456.g004>




Fig 5. The percentage of households where there are 0, 1, 2, 3, 4, 5+ cases across the different methods of scabies status assignment. The results (median and 2.5% to 97.5% quantiles) are plotted for an exemplar input prevalence percentage between 20–30% across all population sizes.

<https://doi.org/10.1371/journal.pntd.0010456.g005>

In Fig 6A, we show how the efficiency of sampling methods changes in response to different underlying scabies assignment approaches for an exemplar sampling percentage and input prevalence between 20–30%. The school sampling strategy overestimates the prevalence when the assignment of scabies includes an age-specific component. In addition, the household




Fig 6. Observed scabies prevalence in samples selected using different (a) assignment methods and (B) sampling percentages. The results (median and 2.5% to 97.5% quantiles) are plotted for an exemplar input prevalence percentage between 20–30% across all population sizes with a sampling percentage between 20–30%. Red dashed lines represent 20% and 30% prevalence. Additional results with differing input prevalence and differing population sizes are presented in S2, S3 and S4 Figs. In panel B, the highest sampling percentages could not be achieved in the school-based sampling strategy due to insufficient population size in the school aged group.

<https://doi.org/10.1371/journal.pntd.0010456.g006>

sampling strategy introduces higher variance compared to simple random sampling when the assignment of scabies includes a household-specific component, because the households with scabies can be over- or under-selected in the samplings. Across all of the scabies assignment methods combined, the simple random sampling strategy produces the narrowest 95% confidence interval for all sampling percentages (Fig 6B). The dependence of observed prevalence in the samples on the underlying scabies assignment approach remains across different sampling percentages (S1 Fig).

Table 2 shows the percentage of our synthetic population that needed to be sampled to achieve sample prevalence within the window of input prevalence +/- a stated precision level

Table 2. Required sample sizes estimated from simulation results for the simple random, household, and school sampling methods to achieve a given precision, combined across all scabies assignment methods.

Population Size	<i>a priori</i> prevalence (%)	Precision								
		Simple Random Sampling			Household Sampling			School Sampling		
		2%	5%	10%	2%	5%	10%	2%	5%	10%
Small	5–10	60%	15%	5%	90%	40%	15%	X	X	X
Small	>10–20	70%	25%	10%	>90%	50%	20%	X	X	X
Small	>20–30	80%	30%	10%	>90%	60%	25%	X	X	X
Small	>30–40	>90%	40%	15%	>90%	70%	30%	X	X	X
Medium	5–10	40%	10%	3%	70%	25%	10%	X	X	X
Medium	>10–20	50%	15%	3%	80%	40%	10%	X	X	X
Medium	>20–30	70%	20%	5%	90%	40%	15%	X	X	X
Medium	>30–40	>90%	20%	5%	>90%	50%	15%	X	X	X
Large	5–10	25%	5%	3%	60%	15%	5%	X	X	X
Large	>10–20	40%	10%	3%	70%	25%	10%	X	X	X
Large	>20–30	50%	10%	3%	80%	30%	10%	X	X	X
Large	>30–40	>90%	15%	3%	>90%	40%	10%	X	X	X

Small, medium, and large population sizes represent ranges of [500, 1500], (1500, 2500), (2500, 4000], respectively. For the scenarios with X's in school sampling, sampling all school-aged children was insufficient to have 95% confidence that prevalence within the selected precision could be obtained. Results stratified by scabies assignment methods are shown in S1, S2, S3, S4 and S5 Tables.

<https://doi.org/10.1371/journal.pntd.0010456.t002>

in 95% of simulations. This table can be used to estimate required sample sizes in real-world populations, based on the population size, an *a priori* estimate of true population prevalence and a desired precision level. For example, for a small population of between 500 and 1500 individuals and an *a priori* estimate of true prevalence between 10–20%, it is necessary to randomly sample 25% of the population to achieve “true prevalence \pm 5%” in 95 out of 100 samples.

We did not run simulations with a sampling percentage higher than 90%. For the scenarios with “>90%” in Table 2, sampling 90% of the population was insufficient to have 95% confidence that prevalence within the selected precision could be obtained. Therefore, we do not report the sample size required for these scenarios.

Table 2 shows that the required sampling percentage of the population (1) increases when greater precision is needed, (2) increases with a higher *a priori* prevalence, and (3) decreases with a larger population.

Discussion

In this study, we present a method to test efficacy of common sampling strategies for scabies in the context of remote Indigenous communities of Australia. To the best of our knowledge, this study is the first to use a simulation-based approach to test the efficacy of scabies sampling strategies, meeting a critical need identified by the WHO [13]. In this section, we discuss how the performance of sampling strategies depends on our assumptions about the relative importance of household- and age-specific scabies transmission, how our analysis can be used in determining sampling size. Then, we provide the strengths and limitations of our study and our future work.

Our results demonstrate how the performance of sampling methods strongly depends on the underlying drivers of scabies transmission, due to the substantial changes in the distribution of scabies prevalence across population groups depending on how scabies spreads. As the precise drivers of scabies distribution within populations are unknown [11, 14, 15, 18–21], we cannot be sure which of the scabies assignment methods we have used is closest to reality. Therefore, it is important to use a sampling approach that performs well across all the scabies assignment methods. Across all the underlying assumptions about scabies prevalence in household and age groups, the simple random sampling strategy produces the narrowest 95% confidence interval for all sampling percentages. Based on our simulations of scabies in synthetic populations and the use of different sampling strategies, simple random sampling is more efficient than household or school sampling, as it requires smaller sample sizes and, for some combinations of true prevalence, population size and desired precision, is the only method that requires a sample size smaller than 90% of the population. Compared to random sampling, household sampling requires larger sample sizes to achieve a desired precision. School-based sampling may result in biased estimates of prevalence due to high prevalence of scabies in school-aged children.

When the aim of undertaking a prevalence survey is only to determine whether prevalence is above or below a threshold, then depending on the *a priori* prevalence assumption, high levels of precision may not be required and a smaller sample size may be sufficient. In such cases Table 2 can be used. Where the desired aim of the sample is to determine whether prevalence is above or below a given threshold, say 10%, for the purposes of running a community treatment day: for example, with a medium size population and an *a priori* estimate of true population prevalence between 20–30%, 10% precision would be sufficient to conclude whether the prevalence is higher than 10%. Adopting a simple random sampling strategy, a sample of 5%

of the population would be sufficient to reach a decision about whether a community treatment day is required. Note that it is better to overestimate *a priori* prevalence than underestimate, as an underestimate could result in an inadequate sample size.

Simulation approaches allow many scenarios to be investigated [29–35]. In this study, we use a range of population sizes and test assignment methods consistent with the literature [11, 14, 15, 18–21]. Our methodology can be applied to test efficacy of sampling strategies for estimating point prevalence of various infectious diseases. We present a pseudo code as a generic framework to compare sampling strategies in measuring disease prevalence in communities (S5 Fig).

In this work, we do not account for the practicality or cost-effectiveness of undertaking the different types of sampling [3]. In addition, we only consider remote Indigenous communities of Australia with population sizes ranging between 500 and 4000 [36]. Further analyses would be useful to estimate effectiveness of sampling strategies in urban or peri-urban areas with larger population sizes [e.g. 18].

In intervention studies, estimation of prevalence pre- and post-intervention may be necessary [10, 20, 40, 41]. In such cases, a period estimation of disease prevalence, therefore a dynamic transmission model, is needed to accurately represent the impact of intervention on the prevalence [29]. The current version of our approach is not applicable for prevalence estimation in these cases since we only create a snapshot of disease prevalence and test the strategies for estimating point prevalence. As future work, the approach proposed here could be extended to consider sampling populations in pre- and post-intervention periods. In addition, our results show that scabies distribution in communities can provide us some clues about the underlying transmission mechanisms (Figs 4 and 5). Our approach can also be extended by comparing these distributions of scabies prevalence in sub-populations to existing survey data of scabies prevalence to infer transmission mechanisms in various populations.

Even though feasibility and cost-effectiveness of sampling strategies are crucial [3], the design of such strategies should take into account the inherent biases that may exist [13, 30, 42]. Due to its feasibility, the WHO recommends school-based sampling to estimate scabies prevalence [13]. However, our results show that the scabies prevalence estimated by using school-based sampling may not be generalisable across the whole community. Our findings highlight the importance of simulation approaches in evaluating and comparing sampling strategies in different population and disease settings.

Supporting information

S1 Fig. Observed scabies prevalence in samples for given sampling percentages, across each of the different scabies assignment methods. The results (median and 2.5% to 97.5% quantiles) are plotted for an exemplar input prevalence percentage between 20–30% across all population sizes where (a) random, (b) high household-specific, (c) mild household-specific, (d) age-specific, (e) age-and-household-specific scabies assignment method is used. Red dashed lines represent 20% and 30% prevalence. Error bars represent the 2.5% to 97.5% quantiles. (TIF)

S2 Fig. Observed scabies prevalence in samples selected using different scabies assignment methods and different input prevalence percentages. The results (median and 2.5% to 97.5% quantiles) are plotted for four exemplar input prevalence percentages of (a) 5%, (b) 10%, (c) 20%, (d) 40% across all population sizes with a sampling percentage of 20%. Red dashed lines represent the input prevalences. (TIF)

S3 Fig. Observed scabies prevalence in samples selected using different sampling methods and sampling percentages across populations with (a) small ([500, 1500]), (b) medium ((1500, 2500)), and (c) large sizes ((2500,4000)). The results (median and 2.5% to 97.5% quantiles) are plotted for an exemplar input prevalence percentage between 20–30% with a sampling percentage of 20%. Red dashed lines represent 20% and 30% prevalence. (TIF)

S4 Fig. Observed scabies prevalence in samples selected using different sampling methods, sampling percentages, and input prevalence. The results (median and 2.5% to 97.5% quantiles) are plotted for four exemplar input prevalence percentages of (a) 5%, (b) 10%, (c) 20%, (d) 40% across all population sizes with a sampling percentage of 20%. Red dashed lines represent the input prevalences. In the school-based sampling strategy the highest sampling percentages could not be achieved due to insufficient population size in the school aged group. (TIF)

S5 Fig. A generic pseudo code for measuring the efficacy of sampling methods in estimating point prevalence of a given disease. (TIF)

S1 Table. Required sample sizes estimated from simulation results for the simple random, household, and school sampling methods to achieve a given precision, where input scabies prevalence is distributed according to the random method. Small, medium, and large population sizes represent ranges of [500, 1500], (1500, 2500], (2500, 4000]. For the scenarios with X's in school sampling, sampling all school-aged children was insufficient to have 95% confidence that prevalence within the selected precision could be obtained. (XLSX)

S2 Table. Required sample sizes estimated from simulation results for the simple random, household, and school sampling methods to achieve a given precision, where input scabies prevalence is distributed according to the high household-specific method. Small, medium, and large population sizes represent ranges of [500, 1500], (1500, 2500], (2500, 4000]. For the scenarios with X's in school sampling, sampling all school-aged children was insufficient to have 95% confidence that prevalence within the selected precision could be obtained. (XLSX)

S3 Table. Required sample sizes estimated from simulation results for the simple random, household, and school sampling methods to achieve a given precision, where input scabies prevalence is distributed according to the mild household-specific method. Small, medium, and large population sizes represent ranges of [500, 1500], (1500, 2500], (2500, 4000]. For the scenarios with X's in school sampling, sampling all school-aged children was insufficient to have 95% confidence that prevalence within the selected precision could be obtained. (XLSX)

S4 Table. Required sample sizes estimated from simulation results for the simple random, household, and school sampling methods to achieve a given precision, where input scabies prevalence is distributed according to the age-specific method. Small, medium, and large population sizes represent ranges of [500, 1500], (1500, 2500], (2500, 4000]. For the scenarios with X's in school sampling, sampling all school-aged children was insufficient to have 95% confidence that prevalence within the selected precision could be obtained. (XLSX)

S5 Table. Required sample sizes estimated from simulation results for the simple random, household, and school sampling methods to achieve a given precision, where input scabies prevalence is distributed according to the age-and-household-specific method. Small, medium, and large population sizes represent ranges of [500, 1500], (1500, 2500], (2500, 4000]. For the scenarios with X's in school sampling, sampling all school-aged children was insufficient to have 95% confidence that prevalence within the selected precision could be obtained. (XLSX)

Acknowledgments

The household composition data used to supplement Australian Bureau of Statistics data in this paper were collected as part of the Life Course Program. We thank the dedicated Life Course research team who traced participants and collected the data. We especially thank the young adults belonging to the Aboriginal Birth Cohort and their families and community for their co-operation and support and all the individuals who helped in the urban and remote locations.

Author Contributions

Conceptualization: Rebecca H. Chisholm, Jodie McVernon, Nicholas Geard, Patricia Therese Campbell.

Formal analysis: Nefel Tellioglu.

Methodology: Nefel Tellioglu.

Supervision: Rebecca H. Chisholm, Nicholas Geard, Patricia Therese Campbell.

Writing – original draft: Nefel Tellioglu.

Writing – review & editing: Rebecca H. Chisholm, Jodie McVernon, Nicholas Geard, Patricia Therese Campbell.

References

1. Fuller LC. Epidemiology of scabies. *Current Opinion in Infectious Diseases*. 2013; 26(2):123–126. <https://doi.org/10.1097/QCO.0b013e32835eb851> PMID: 23411418
2. DALYs and HALE Collaborators. Global, regional, and national disability-adjusted life-years (DALYs) for 333 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet*. 2017; 390(10100):1260–1344. [https://doi.org/10.1016/S0140-6736\(17\)32130-X](https://doi.org/10.1016/S0140-6736(17)32130-X)
3. Engelman D, Steer AC. Control strategies for scabies. *Tropical Medicine and Infectious Disease*. 2018; 3(3):98. <https://doi.org/10.3390/tropicalmed3030098> PMID: 30274494
4. Engelman D, Cantey PT, Marks M, Solomon AW, Chang AY, Chosidow O, et al. The public health control of scabies: priorities for research and action. *The Lancet*. 2019; 394(10192):81–92. [https://doi.org/10.1016/S0140-6736\(19\)31136-5](https://doi.org/10.1016/S0140-6736(19)31136-5) PMID: 31178154
5. Tsoi SK, Lake SJ, Thean LJ, Matthews A, Sokana O, Kama M, et al. Estimation of scabies prevalence using simplified criteria and mapping procedures in three Pacific and southeast Asian countries. *BMC Public Health*. 2021; 21(1):2060. <https://doi.org/10.1186/s12889-021-12039-2> PMID: 34758806
6. Gibbs S. Skin disease and socioeconomic conditions in rural Africa: Tanzania. *International Journal of Dermatology*. 1996; 35(9):633–639. <https://doi.org/10.1111/j.1365-4362.1996.tb03687.x> PMID: 8876289
7. Ferrieri P, Dajani AS, Wannamaker LW, Chapman SS, et al. Natural history of impetigo: I. Site sequence of acquisition and familial patterns of spread of cutaneous streptococci. *The Journal of Clinical Investigation*. 1972; 51(11):2851–2862. <https://doi.org/10.1172/JCI107108> PMID: 5080412
8. Currie BJ, Carapetis JR. Skin infections and infestations in Aboriginal communities in northern Australia. *Australasian Journal of Dermatology*. 2000; 41(3):139–143. <https://doi.org/10.1046/j.1440-0960.2000.00417.x> PMID: 10954983

9. Wong LCF, Amega B, Connors C, Barker R, Dulla ME, Currie BJ. Outcome of an interventional program for scabies in an Indigenous community. *Medical Journal of Australia*. 2001; 175(7):367–370. <https://doi.org/10.5694/j.1326-5377.2001.tb143620.x> PMID: 11700814
10. Andrews RM, Kearns T, Connors C, Parker C, Carville K, Currie BJ, et al. A regional initiative to reduce skin infections amongst aboriginal children living in remote communities of the Northern Territory, Australia. *PLoS Neglected Tropical Diseases*. 2009; 3(11):e554. <https://doi.org/10.1371/journal.pntd.0000554> PMID: 19936297
11. Romani L, Steer AC, Whitfield MJ, Kaldor JM. Prevalence of scabies and impetigo worldwide: a systematic review. *The Lancet Infectious Diseases*. 2015; 15(8):960–967. [https://doi.org/10.1016/S1473-3099\(15\)00132-2](https://doi.org/10.1016/S1473-3099(15)00132-2) PMID: 26088526
12. Reid HF, Birju B, Holder Y, Hospedales J, Poon-King T. Epidemic scabies in four Caribbean islands, 1981–1988. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 1990; 84(2):298–300. [https://doi.org/10.1016/0035-9203\(90\)90296-Q](https://doi.org/10.1016/0035-9203(90)90296-Q) PMID: 2389327
13. Engelman D, Marks M, Steer AC, Beshah A, Biswas G, Chosidow O, et al. A framework for scabies control. *PLoS Neglected Tropical Diseases*. 2021; 15(9):e0009661. <https://doi.org/10.1371/journal.pntd.0009661> PMID: 34473725
14. Hay R, Steer A, Engelman D, Walton S. Scabies in the developing world—its prevalence, complications, and management. *Clinical Microbiology and Infection*. 2012; 18(4):313–323. <https://doi.org/10.1111/j.1469-0691.2012.03798.x> PMID: 22429456
15. Dagne H, Dessie A, Destaw B, Yallew WW, Gizaw Z. Prevalence and associated factors of scabies among schoolchildren in Dabat district, northwest Ethiopia, 2018. *Environmental Health and Preventive Medicine*. 2019; 24:67. <https://doi.org/10.1186/s12199-019-0824-6> PMID: 31785612
16. Walton SF, Choy JL, Bonson A, Valle A, McBroom J, Taplin D, et al. Genetically distinct dog-derived and human-derived *Sarcoptes scabiei* in scabies-endemic communities in northern Australia. *The American Journal of Tropical Medicine and Hygiene*. 1999; 61(4):542–547. <https://doi.org/10.4269/ajtmh.1999.61.542> PMID: 10548286
17. Walton SF, Dougall A, Pizzutto S, Holt D, Taplin D, Arlian L, et al. Genetic epidemiology of *Sarcoptes scabiei* (Acari: Sarcoptidae) in northern Australia. *International Journal for Parasitology*. 2004; 34(7):839–849. <https://doi.org/10.1016/j.ijpara.2004.04.002> PMID: 15157767
18. Collinson S, Timothy J, Zayzay SK, Kollie KK, Lebas E, Candy N, et al. The prevalence of scabies in Monrovia, Liberia: a population-based survey. *PLoS Neglected Tropical Diseases*. 2020; 14(12):e0008943. <https://doi.org/10.1371/journal.pntd.0008943> PMID: 33284821
19. Enbiale W, Ayalew A. Investigation of a scabies outbreak in drought-affected areas in Ethiopia. *Tropical Medicine and Infectious Disease*. 2018; 3(4):114. <https://doi.org/10.3390/tropicalmed3040114> PMID: 30380650
20. Kearns TM, Speare R, Cheng AC, McCarthy J, Carapetis JR, Holt DC, et al. Impact of an ivermectin mass drug administration on scabies prevalence in a remote Australian Aboriginal community. *PLoS Neglected Tropical Diseases*. 2015; 9(10):e0004151. <https://doi.org/10.1371/journal.pntd.0004151> PMID: 26516764
21. Romani L, Koroivueta J, Steer AC, Kama M, Kaldor JM, Wand H, et al. Scabies and impetigo prevalence and risk factors in Fiji: a national survey. *PLoS Neglected Tropical Diseases*. 2015; 9(3):e0003452. <https://doi.org/10.1371/journal.pntd.0003452> PMID: 25738499
22. Carapetis JR, Connors C, Yarmirr D, Krause V, Currie BJ. Success of a scabies control program in an Australian Aboriginal community. *The Pediatric Infectious Disease Journal*. 1997; 16(5):494–499. <https://doi.org/10.1097/00006454-199705000-00008> PMID: 9154544
23. Eason R, Tasman-Jones T. Resurgent yaws and other skin diseases in the Western Province of the Solomon Islands. *Papua and New Guinea Medical Journal*. 1985; 28(4):247–250. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/2940770>. PMID: 2940770
24. Lake SJ, Engelman D, Sokana O, Nasi T, Boara D, Grobler AC, et al. Defining the need for public health control of scabies in Solomon Islands. *PLoS Neglected Tropical Diseases*. 2021; 15(2):e0009142. <https://doi.org/10.1371/journal.pntd.0009142> PMID: 33617544
25. Thean LJ, Romani L, Engelman D, Jenney A, Wand H, Mani J, et al. Prospective Surveillance of Primary Healthcare Presentations for Scabies and Bacterial Skin Infections in Fiji, 2018–2019. *The American Journal of Tropical Medicine and Hygiene*. 2021; 105(1):230–237. <https://doi.org/10.4269/ajtmh.20-1459> PMID: 34029210
26. Matthews A, Le B, Amaral S, Arkell P, Monteiro M, Clarke N, et al. Prevalence of scabies and impetigo in school-age children in Timor-Leste. *Parasites & Vectors*. 2021; 14:156. <https://doi.org/10.1186/s13071-021-04645-1> PMID: 33722285
27. La Vincente S, Kearns T, Connors C, Cameron S, Carapetis J, Andrews R. Community management of endemic scabies in remote aboriginal communities of northern Australia: low treatment uptake and high

- ongoing acquisition. *PLoS Neglected Tropical Diseases*. 2009; 3(5):e444. <https://doi.org/10.1371/journal.pntd.0000444> PMID: 19478832
28. Zayyid MM, Saadah RS, Adil A, Rohela M, Jamaiah I. Prevalence of scabies and head lice among children in a welfare home in Pulau Pinang, Malaysia. *Tropical Biomedicine*. 2010; 27(3):442–446. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21399584>.
 29. Giardina F, Coffeng LE, Farrell SH, Vegvari C, Werkman M, Truscott JE, et al. Sampling strategies for monitoring and evaluation of morbidity targets for soil-transmitted helminths. *PLoS Neglected Tropical Diseases*. 2019; 13(6):e0007514. <https://doi.org/10.1371/journal.pntd.0007514> PMID: 31242194
 30. Schmidt WP, Genser B, Barreto ML, Clasen T, Luby SP, Cairncross S, et al. Sampling strategies to measure the prevalence of common recurrent infections in longitudinal studies. *Emerging Themes in Epidemiology*. 2010; 7:5. <https://doi.org/10.1186/1742-7622-7-5> PMID: 20678239
 31. Fronterre C, Amoah B, Giorgi E, Stanton MC, Diggle PJ. Design and analysis of elimination surveys for neglected tropical diseases. *The Journal of Infectious Diseases*. 2020; 221(Supplement_5):S554–S560. <https://doi.org/10.1093/infdis/jiz554> PMID: 31930383
 32. Alexander CJ, Holland JM, Winder L, Woolley C, Perry JN. Performance of sampling strategies in the presence of known spatial patterns. *Annals of Applied Biology*. 2005; 146(3):361–370. <https://doi.org/10.1111/j.1744-7348.2005.040129.x>
 33. Toor J, Coffeng LE, Hamley JI, Fronterre C, Prada JM, Castaño MS, et al. When, who, and how to sample: designing practical surveillance for 7 neglected tropical diseases as we approach elimination. *The Journal of Infectious Diseases*. 2020; 221(Supplement_5):S499–S502. <https://doi.org/10.1093/infdis/jiaa198> PMID: 32529261
 34. Diggle PJ, Amoah B, Fronterre C, Giorgi E, Johnson O. Rethinking neglected tropical disease prevalence survey design and analysis: a geospatial paradigm. *Transactions of The Royal Society of Tropical Medicine and Hygiene*. 2021; 115(3):208–210. <https://doi.org/10.1093/trstmh/tra020> PMID: 33587142
 35. Fornace KM, Fronterre C, Fleming FM, Simpson H, Zoure H, Rebollo M, et al. Evaluating survey designs for targeting preventive chemotherapy against *Schistosoma haematobium* and *Schistosoma mansoni* across sub-Saharan Africa: a geostatistical analysis and modelling study. *Parasites & Vectors*. 2020; 13:555. <https://doi.org/10.1186/s13071-020-04413-7> PMID: 33203463
 36. Dockery AM, Singh R, Harris M, Holyoak N. Projecting Aboriginal and Torres Strait Islander populations for remote communities: a small number approach; 2017. [Cited 30 September 2021]. Available from: https://nintione.com.au/resource/CW030_ProjectingAboriginalTorresStraitIslanderPopulations_ASmallNumberApproach.pdf.
 37. Australian Bureau of Statistics. SA2-112 Household & Family Composition by Number of Persons Usually Resident by Indigenous Status of Household-Census 2016; 2017. [Cited 30 September 2021]. Available from: <https://data.gov.au/dataset/ds-aurin-00dabd52f6c09f7620699bf812bb2351cbca715b941a88a89752fee2d3f18409/details?q=>.
 38. Vino T, Singh GR, Davison B, Campbell PT, Lydeamore MJ, Robinson A, et al. Indigenous Australian household structure: a simple data collection tool and implications for close contact transmission of communicable diseases. *PeerJ*. 2017; 5:e3958. <https://doi.org/10.7717/peerj.3958> PMID: 29085755
 39. Australian Health Ministers' Advisory Council. Aboriginal and Torres Strait Islander Health Performance Framework 2017; 2017. [Cited 25 May 2021]. Available from: https://www.niaa.gov.au/sites/default/files/publications/2017-health-performance-framework-report_1.pdf.
 40. Matthewman J, Manego RZ, Dimessa Mbadinga LB, Šinkovec H, Völker K, Akinosho M, et al. A randomized controlled trial comparing the effectiveness of individual versus household treatment for Scabies in Lambaréné, Gabon. *PLoS Neglected Tropical Diseases*. 2020; 14(6):e0008423. <https://doi.org/10.1371/journal.pntd.0008423> PMID: 32589632
 41. Romani L, Marks M, Sokana O, Nasi T, Kamoriki B, Cordell B, et al. Efficacy of mass drug administration with ivermectin for control of scabies and impetigo, with coadministration of azithromycin: a single-arm community intervention trial. *The Lancet Infectious Diseases*. 2019; 19(5):510–518. [https://doi.org/10.1016/S1473-3099\(18\)30790-4](https://doi.org/10.1016/S1473-3099(18)30790-4) PMID: 30956111
 42. Alonzo TA, Pepe MS, Lumley T. Estimating disease prevalence in two-phase studies. *Biostatistics*. 2003; 4(2):313–326. <https://doi.org/10.1093/biostatistics/4.2.313> PMID: 12925524