

Mobile Health interventions for modifying indigenous maternal and child-health related behaviours: a systematic review

Sana Ishaque, Ola Ela, Anna Dowling, Chris Rissel, Karla Canuto, Kerry Hall, Niranjan Bidargaddi, Annette Briley, Claire T Roberts, Billie Bonevski

Submitted to: Journal of Medical Internet Research
on: February 01, 2024

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript.....	5
Supplementary Files.....	27
0.....	27
0.....	27
.....	27
CONSORT (or other) checklists.....	28
CONSORT (or other) checklist 0.....	28



Mobile Health interventions for modifying indigenous maternal and child-health related behaviours: a systematic review

Sana Ishaque^{1*}; Ola Ela¹; Anna Dowling^{1*}; Chris Rissel¹; Karla Canuto^{1*}; Kerry Hall^{2*}; Niranjana Bidargaddi^{1*}; Annette Briley^{1*}; Claire T Roberts^{1*}; Billie Bonevski^{1*}

¹ Flinders University College of Medicine and Public Health Bedford Park AU

² Griffith University Queensland AU

*these authors contributed equally

Corresponding Author:

Sana Ishaque

Flinders University
College of Medicine and Public Health
Sturt Road
Bedford Park
AU

Abstract

Background: The transition to motherhood is a pivotal time for promoting healthy behaviours, particularly among indigenous women, who encounter significant barriers to accessing health information. Mobile health interventions (mHealth) promoting healthy lifestyle changes, offer an adaptable and inexpensive method for improving access health information but require cultural appropriateness and suitability for acceptance and effectiveness in indigenous populations. No systematic review on effective mHealth interventions for indigenous women during pregnancy and the early childhood years has been conducted.

Objective: This study evaluated the effectiveness of mHealth interventions, promoting healthy lifestyle changes, for indigenous mothers and children from conception to five years post-partum. It explored the effectiveness differences based on participant engagement, intervention nature, and provision of context.

Methods: A systematic search of five databases; SCOPUS, MEDLINE, CINAHL, PsycINFO, and ProQuest (Dissertation or Thesis); was conducted to identify studies focusing on maternal and child health, indigenous populations, and mHealth following a pre-registered PROSPERO protocol (CRD42023395710). HealthInfoNet was searched for grey literature and the reference lists of included studies were hand searched. Randomised controlled trials and other interventional study designs including pre-post comparison and cohort studies were eligible for inclusion. Quality of studies was evaluated by two independent reviewers using the Mixed Methods Quality Appraisal Tool and the Centre of Excellence in Aboriginal Chronic Disease Knowledge Translation and Exchange tool. Details from the studies was extracted using a predeveloped extraction form and a descriptive synthesis of the data was performed.

Results: Of the 688 articles screened, only three met the eligibility criteria. Each paper evaluated a different mHealth intervention. The three mHealth interventions identified were: Remote Prenatal Education, the Short Messaging System Parent Action Intervention (two-way text messaging), and the Screening, Brief Intervention and Referral to Treatment eCHECKUP To Go (web-based screening and intervention).

While all studies reported significant outcomes for varying parts of their respective interventions, they all had a small sample size, and none provided a rationale for the power calculation of the sample size for the outcomes reported. Therefore, it was not possible to establish whether the differences in the effectiveness were due to the interventions.

Conclusions: The current literature does not have any evidence of the effectiveness of mHealth interventions for maternal and child health behaviour change. A thorough evaluation with consideration of cultural contexts and user preferences during intervention design and development is crucial for maximizing their potential. Despite scant evidence, mHealth interventions, hold promise for enhancing indigenous mothers' health.

(JMIR Preprints 01/02/2024:57019)

DOI: <https://doi.org/10.2196/preprints.57019>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in [JMIR Publications](#).

No. Please do not make my accepted manuscript PDF available to anyone.

Original Manuscript



Mobile Health interventions for modifying indigenous maternal and child-health related behaviours: a systematic review

Abstract

Background

The transition to motherhood is a pivotal time for promoting healthy behaviours, particularly among indigenous women, who encounter significant barriers to accessing health information. Mobile health interventions (mHealth) promoting healthy lifestyle changes, offer an adaptable and inexpensive method for accessing health information but require cultural appropriateness and suitability for acceptance and effectiveness in indigenous populations. No systematic review on effective mHealth interventions for indigenous women during pregnancy and the early childhood years has been conducted.

Objectives

This review evaluated the effectiveness of mHealth interventions, promoting healthy behaviours, for indigenous mothers and children from conception to five years post-partum. It also aimed to explore the observed effectiveness differences based on participant engagement, intervention design and provision of context. Further, the review explored if the interventions were co-designed.

Methods

A systematic search of five databases; SCOPUS, MEDLINE, CINAHL, PsycINFO, and ProQuest (Dissertation or Thesis) was conducted. Studies were included if they were either a randomised controlled trial, pre-post comparison, or a cohort study using mHealth with indigenous women for maternal and child health following a pre-registered PROSPERO protocol (CRD42023395710). HealthInfoNet was searched for grey literature and the reference lists of included studies were hand searched. Initial title and abstract screen for eligibility was performed by one reviewer. Full text screen of eligible studies and quality appraisal of included studies was performed by two reviewers independently. The appraisal tools used were Mixed Methods Quality Appraisal Tool and the Centre of Excellence in Aboriginal Chronic Disease Knowledge Translation and Exchange. A descriptive synthesis of the extracted data was performed.

Results

Of the 663 articles screened, only three met the eligibility criteria. Each paper evaluated a different mHealth intervention. The three mHealth interventions identified were: Remote Prenatal Education; the Short Messaging System Parent Action Intervention (two-way text messaging); and the Screening, Brief Intervention and Referral to Treatment eCHECKUP To Go (web-based screening and intervention).

Statistically significant changes were reported in some outcomes including increase in parental participation rate in a face-to-face prenatal education; increased rate of breastfeeding initiation, and exclusive breastfeeding (2-12 months); improved overall children's behaviour related to sleep, diet, physical activity, screen time, and intake of sugary beverages; improved individual child behaviours related to physical activity and sleep; and decrease in alcohol drinks/week and binge drinking episodes/2 weeks due to time effect. However, no study provided a sample size calculation for the reported significant outcomes.

Also, due to the small number of included studies and each study evaluating a different intervention, it was not possible to combine results to ascertain if the participant engagement, intervention design or community context had any impact on the effectiveness.

Conclusion

Due to the lack of sample size calculation, it was not possible to establish whether differences in the effectiveness were due to the interventions or a Type 1 statistical error. Therefore, caution is required in the interpretation of these findings.

Abbreviations

- CREATE – the Centre of Excellence in Aboriginal Chronic Disease Knowledge Translation and Exchange
- mHealth – Mobile health technology/intervention
- PRISMA – Preferred Reporting Items for Systematic Reviews and Meta-Analyses
- PROSPERO – Prospectively registered systematic reviews
- RCT – randomised controlled trial

Keywords

Indigenous, co-design, maternal, children, digital health, mobile health.

Introduction

Globally, indigenous populations share experiences of devastating disruptions to development due to violent settler colonial practices which persisted well into the twentieth century and continue to impact today[1]. This historical suffering and the consequent vestiges are inextricably linked to current overrepresentation of indigenous people in population statistics relating to avoidable mortality, disease burden, and social and economic disadvantage[2]. To note, in this review, the term indigenous is used for indigenous populations across the globe.

Adverse health outcomes including low birth weight, preterm birth, stillbirths, and perinatal mortality rates disproportionately affect indigenous families [3, 4]. Indigenous women are more likely to experience domestic violence, have high levels of psychological distress and exhibit a higher prevalence of behavioural risk factors including tobacco, alcohol and other drug use in comparison with non-indigenous women[5-7]. Disparities between indigenous and non-indigenous maternal and child health outcomes highlight the need for urgent and specialised support [3]. The perinatal period offers a unique opportunity for women to adopt healthy behaviours for the benefit of both themselves and their baby [8]. In addition, experiences during the first 2000 days of a child's life have a significant lasting impact on cognitive, physical, social, and emotional health[9, 10]. Therefore, interventions applied during the perinatal period and early childhood are more successful than those implemented at a later stage of life[9].

Of relevance is the geographic impact on health inequities experienced by indigenous populations. Compared with urban or regional populations, communities in rural and remote regions experience higher mortality [11]. The disparity is attributed to barriers associated with high cost, poor access and culturally inappropriate service provision, as well as employment, education, and income disadvantages [12].

Mobile Health interventions offer an alternative option for the provision and dissemination of health information, particularly for communities outside major cities where health services reach is reduced. Health service providers are increasingly using digital technology for health promotion due to its practicality, reach and accessibility [13]. mHealth is being used for education, behaviour modification, data collection and tracking, point of care diagnostics, health decision support and record keeping [14, 15]. This coincides with the increased use of mobile technology and specifically, its utilisation for health and social and emotional wellbeing (SEWB) purposes by indigenous women

[16, 17].

It has been established that indigenous women are likely to successfully engage in services during the antenatal period when they feel empowered through education, family/community support and cultural connectedness[4, 6, 18, 19]. Effective health promotion are accessible, empowering, respect the values of indigenous family structures and kinship systems and most critically, are indigenous-led [20]. Like any other intervention, mHealth when used for indigenous populations requires the intervention to be culturally appropriate and acceptable[21].

There are very few studies that have reported the benefits of culturally responsive mHealth interventions specifically developed to support the healthy lifestyle choices for indigenous women[22-24]. Research that fails to consult with indigenous communities has shown to provide little benefit [10]. The homogenisation of applying mainstream health interventions risks perpetuating inequities, hindering engagement and discounting the legacy of historical atrocities [25]. It is critical that mHealth are culturally safe and acceptable and their evaluations consider differences in participant characteristics, features of the intervention and the context of delivery [26]. The primary objective of this systematic review was to assess the effectiveness of mHealth interventions designed for use by indigenous mothers of children (conception to five years postpartum) that seek to promote healthy behaviours. The secondary aims of this review included examining the observed differences in the effectiveness of the interventions based on participant engagement, the intervention design and the programmatic and community context in which the intervention is implemented. This review also explored if the mHealth intervention has been designed or adapted specifically for indigenous populations.

Methods

A systematic review protocol outlining the search strategy, methods for title/abstract and full-text screening, data extraction template, quality assessment methods, data analysis methods, synthesis and statistical issues, publication bias and any conflicts of interest was developed and registered with the international database of prospectively registered systematic reviews (PROSPERO), registration number: (CRD42023395710) [27]. The review followed Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guideline for the conduct and reporting [28]. The protocol was followed fully except that the third aim to explore if the identified interventions were co-designed was included after the protocol was submitted for registration to PROSPERO based on experts' advice.

Eligibility criteria

A publication was eligible for inclusion if it reported findings from a randomised controlled trial (RCT), pseudorandomised controlled trial, comparative study with concurrent controls, or pre-post study that applied a user operated digital health intervention as an intervention. Population included was self-identified indigenous women in the antenatal period and/or with children aged 0 to 5 years of age. The outcomes eligible for inclusion were health behaviours during pregnancy, which related to diet, use of tobacco, alcohol, other drugs, physical activity; and behaviours performed by mothers and/or other caregivers during the early childhood period, including behaviours related to health and hygiene care, feeding, stimulation, responsiveness, and safety. The review was restricted to publications reported in English. There was no restriction on the type of control, type of digital health intervention, country, setting in which the study was conducted, nor date of publication.

Studies were excluded if they reported on the development or design of a digital health intervention without reporting on its effectiveness on the behavioural outcomes of indigenous women/children, reported on mixed indigenous and non-indigenous populations without reporting subgroup data, or applied the intervention directly to children with the intention to change maternal behaviour.

In this review, user-operated digital health technologies are regarded as any form of computerised

technology that the intended (beneficiary or target) audience interacts with in a way that is not mediated by a third-party (e.g., by a healthcare provider). This includes mobile/cellular phone technologies (or mHealth); web 2.0 technologies (inclusive of social media); websites and web-based applications; and non-web applications (e.g., delivered via offline electronic devices).

Search Strategy

With the assistance of a research librarian, a systematic search was performed in SCOPUS, MEDLINE, CINAHL, PsycINFO, and ProQuest (Dissertation or Thesis) in Oct 2022 and was updated in Dec 2023. The search was conducted using controlled vocabulary and keywords related to the terms: maternal, child/family health, indigenous populations, and mobile health. Specially designed search filters have been used to identify terms for indigenous populations [29]. Details of the search strategy for all databases is given in Appendix 1. The search strategy was modified to adapt to variations in indexing among the other databases and complete search strategy is available on request. Additionally, HealthInfoNet website and reference lists of the included studies were hand searched for relevant grey literature and any additional relevant studies respectively.

Title/abstract and full-text screening

The literature screening was performed using the Covidence systematic review software. After removing duplicate articles, title and abstract screen was performed by one reviewer per article (SI, OE, AD). The studies selected for full text review were screened by two independent assessors (SI, OE, AD) against the eligibility criteria, with conflicts managed with discussion and with the assistance of an expert reviewer (BB).

Data Extraction and Analysis

Data extraction was performed by one assessor (either OE or AD) using a template designed to extract information about the study methodology, population characteristics (including self-identification of indigenous status), behavioural target, intervention characteristics, intervention context, information about co-design of the intervention and reported outcomes. Another reviewer (SI) checked the extracted data for accuracy and completeness.

The data is presented descriptively and in narrative form, where applicable. The planned analysis to perform meta-aggregation of qualitative data and meta-analysis of quantitative data was not feasible due to the small number of included studies.

Risk of Bias Assessment

An assessment of the risk of bias for included studies was performed independently by two reviewers (OE, AD) using the Mixed Methods Appraisal Tool [30] and the Centre of Excellence in Aboriginal Chronic Disease Knowledge Translation and Exchange (CREATE) tool [31]; with a third assessor (SI) managing conflicts. Anticipating a relatively small body of literature, risk of bias assessment was not used to exclude otherwise eligible literature but is presented and discussed alongside the review results.

Results


Eligibility


From the database search 867 articles were retrieved, of which 229 were duplicates. After removal of duplicates, title and abstract of 663 articles were screened for relevance, of which 42 were found eligible for full text screening. After full-text screening of 42 articles, three met the inclusion criteria and were included in this systematic review (Fig. 1) [39-41]. There were n=39 excluded studies due

to wrong outcomes (5 studies), wrong intervention (6 studies), wrong study design (12 studies), wrong patient population (13 studies), wrong publication – book (1 study), and protocol only - no results (2 studies).

An additional six articles were found eligible for full text screen on hand search of reference lists of the included studies, none of them were included in the review as the study design was inappropriate for this review.

Figure. 1 PRISMA flow diagram





Preprint
JMIR Publications

Hui et al. [40]	To assess the impact of community-based remote prenatal education	Sagkeeng and Sandy Bay (two rural Ojibwe First Nation communities) Garden Hill (a remote Anishinew First Nation community) Internet available at all locations; Speed of Wifi at Garden Hill unable to transfer high quality video images	Canada	Community based	Pre-post comparison	42% attendance (n=53) in the communities for the social media assisted prenatal chat group	Pregnant women Total number of participants not reported	Breastfeeding initiation, Breastfeeding duration, Parental participation in parental education program	Outcomes 1-year pre and post intervention : Participation rate in prenatal program across the communities increased significantly 36.4% (85/233 births) V 54.5% (125/231 births) $P<.001$. Significant increase in breastfeeding initiation rate in only 1/3 of the communities 23% (n=15/65 live births) to 67% (n=288/42 live births) $P<.001$, No significant difference in breastfeeding initiation rate in Sandy Bay and Garden Hill areas.
-----------------	---	---	--------	-----------------	---------------------	--	---	--	---

Brown et al. [39]	This study investigated the feasibility of delivering health-related short text messages to parents to reduce obesity risk among children aged 3 to 5 years	American Indian reservation	USA	Community based	Pre-post comparison	17/17 completed 5 weeks study	Parents of children 3 to 5 years of age American Indian (47%), mean age 34 years, female (15/17, 88%), annual incomes greater than \$30,000 (59%), college graduates (71%) Total n = 17	Child behaviours related to sleep, diet, physical activity, screen time, and intake of sugary beverages	Outcome compared pre and post intervention: Significant change in favour of the intervention in overall child behaviour (P=.051) Significant difference in child behaviour related to physical activity and sleep (P=.046) No significant difference in child behaviour related to diet
Montag et al. [41]	To develop and test an adaptation of an online intervention for reducing risky drinking in Native women of childbearing	American Indian/Alaska Native women	USA	Community based	Randomised controlled trial	263 recruited, 16 (6.1%) lost to follow up	Sample size for which demographics were reported n = 247 Age: 28.6 _ 0.5 (28.6 _ 0.5) Intervention n = 129 Control n = 134 Total n=263	Binge drinking, Weekly drinking	No significant difference between I and C for alcoholic drinks/week and binge episodes/two weeks Alcoholic drinks/week and binge episodes/two weeks significantly reduced due to time effect P<.05

The publication dates of the three included studies were 2021, 2019, and 2015 (Table 1) [39-41]. Two of the studies were conducted in USA with American First Nation populations [39, 41] and one was conducted in Canada [40] and included three First Nation communities of Canada (Sagkeeng, Sandy Bay, and Garden Hill). The study designs were pre-post comparison (n=2), and randomised controlled trial (n=1) respectively. The health conditions around which behavioural changes were sought included: participation in the prenatal program and breastfeeding initiation [40], childhood obesity [39], and alcohol addiction (or problem drinking)[41]. The outcomes included: breastfeeding initiation, breastfeeding duration, parental participation in parental education program [40],[39] and [41]; child behaviours related to sleep, diet, physical activity, screen time, and intake of sugary beverages [39]; and binge drinking and weekly drinking[41]. The interventions were applied in community settings in all three included studies.

Study Interventions

The eHealth intervention, Remote Prenatal Education, in the pre-post, Hui et al. (2021) was a sub-study of a large intervention that also included some face-to-face components (Table 2). The eHealth component consisted of traditionally/culturally tailored web-accessible educational material on pregnancy, prenatal, and postnatal period related topics such as healthy eating, recipes, traditional foods, food label reading, and nutrition content for common foods and a social media assisted prenatal chat group operated through Facebook messenger. The chat was moderated by community Elders, Canadian Prenatal Nutrition Program (CPNP) workers or the study coordinator. The intervention was accessible via computer/smartphone and could be opened via Facebook. The educational material on the website was available as readings, audios, and videos; news on the

prenatal or postnatal programming in the communities and traditional knowledge regarding a healthy pregnancy was also available. Traditional methods including prayer, sharing circles, and storytelling, were incorporated in breastfeeding education. While the total number of participants was not reported, the participation rate for the social media prenatal group was at 42% (n=53).

The intervention, the Short Messaging System (SmS) Parent Action Intervention, in the pre-post, Brown et al. (2019) study consisted of three short text messages per week for five consecutive weeks sent between 30 Jan to 17 March 2017 for n=17 families (Table 2). The messages were tailored for age and gender of participating parents and children, sent on the phone at individualised times of the day using an online text automation platform, Mosio. The text messages provided information around recommended healthy child nutrition, physical activity, sleep, and screen time behaviours of preschool children. The text message topics were derived from the, 'Let's Go! 5-2-1-0' US national campaign and recommendations for sleep in children.

The intervention in the randomised-control trial utilised the Screening, Brief Intervention, and Referral to Treatment eCHECKUP To Go; which was an online intervention that screened participants for their use of alcohol and based on their answers provided feedback about participants' risk of an alcohol exposed pregnancy and its effect on the foetus (Table 2) [41]. After the initial screening, resources and information on support services were provided to participants. The intervention took approximately 20 minutes to complete. There were 129 women in the intervention group and 134 women in the control group that received 'usual care' which included non-specific educational resources available in waiting areas.

Table 2 Intervention characteristics

Study ID	Intervention Name and details Is the intervention live and any links	Was the digital health intervention being part of a bigger intervention (e.g. there were some face-to-face components)	How was the intervention delivered? (e.g. computer, tablet, phone) and Accessed via	Intervention Context	Name of Indigenous Nation/Country	How many times the participant did interact with the intervention. E.g. Text messages sent thrice a week etc.	Who (if anyone) delivered/moderated the intervention?
Hui et al. 2021 [40]	Remote Prenatal education via an educational website and social media-assisted prenatal chat groups (Facebook or Messenger) Live website (https://momsinmotion.ca/)	Yes	Computer and smartphone Facebook link with website	community-based remote and some face-to-face components	Sagkeeng and Sandy Bay: two rural Ojibwe First Nation communities, Garden Hill: Anishinew First Nation community	Not reported	Community Elders, Canadian Prenatal Nutrition Program workers, or the study coordinator served as the hosts of the Facebook chat groups
Brown et al. 2019 [39]	Two-way text messaging the Short Messaging System Parent Action Intervention The text message topics were derived from the, 'Let's Go! 5-2-1-0' US national campaign and recommendations http://sleepforkids.org/	No	Text messages were sent using Mosio, an online platform for text message automation	Community	American and non-American Indian residing on American Indian Reserves	Three text messages each week for five consecutive weeks	Messages appeared to be sent from a local paediatrician or a tribal health dietitian
Montag et al. 2015 [41]	Screening, Brief Intervention, and Referral to Treatment eCHECKUP TO GO, a web-based brief assessment and intervention tool Participants received individualized web-based	No	Computer, phone, tablet Could be accessed by the website	Participants were recruited from health American Indian/Alaska a Native health clinic	American Indian/Alaska Native women in Southern California	1, 3 and 6 months	Not applicable

	feedback at the end of the session regarding their risk for an AEP, the impact of alcohol exposure to the fetus, the physical and financial cost of their alcohol consumption, and how their drinking compared with that of other Native women. approximately 20 minutes to complete. https://www.echeckuptogo.com/			Intervention delivered in community setting			
--	---	--	--	---	--	--	--

Study Outcomes

In the Hui et al (2021) study, a significant increase was reported in the participation of pregnant women in a face-to-face prenatal program 1 year after the start of intervention compared with pre-program participation rates (Table 1)[40]. Furthermore, a significant increase was reported in the breastfeeding initiation rate at one of the study sites and exclusive breastfeeding between 2 to 12 months was also significantly increased at another study site.

In the Brown (2019) study a pre-post comparison of the five-week text intervention, demonstrated that there was a significant increase in, parent reported, overall child behaviours related to sleep, diet, physical activity, screen time, and intake of sugary beverages in favour of the intervention (Table 1) [39]. When individual behaviours were assessed, there was significant change in child behaviours related to physical activity and sleep and there was no change in child behaviour related to diet.

The Montag (2015) study, which was a randomised controlled trial from April 2011 to Sep 2012, did not find any statistically significant differences in intake of alcoholic drinks per week and binge drinking episodes per week between the intervention and control groups (Table 1) [41].

Cultural Adeptness

The steps and strategies taken to ensure cultural appropriateness of the mHealth interventions were extracted from the included studies, Table 3. Hui et al (2021) and Montag et al (2015) cited previous work that reported the interventions being co-designed for indigenous populations, and Brown et al (2019) used an intervention that was not reported to be co-designed or adapted for indigenous populations [39-41]. In the instances when previous research was cited, those articles were retrieved, and relevant data were extracted in Table 3. Authors of the included studies were also contacted for information related to the co-design/adaptation.

Study ID	Was the intervention culturally adapted (and/or co-designed)?	Is there any reference/citation provided about cultural adaptation/co-design of the intervention?	How was the cultural adaptation done (e.g. design processes/co-design)?	What features were included to make it culturally appropriate?	Were Indigenous people included in the design/adaptation?	What other population(s) were included in the cultural adaptation/co-design process?	Theory (or framework) used for cultural adaptation/co-design?	Evaluation of cultural appropriateness
Hui et al. 2021	Yes	Yes[42]	Not reported	The intervention was built on the previous	Yes	None	Not reported	Not reported

[40]				work of the researchers that identified barriers to participation among the study population [42]				
Brown et al. 2019 [39]	Not reported	No	Not reported	Research approved by the Institutional Review Board at the American Indian Tribal college on the reservation; the design, implementation, and interpretation were done in partnership	Yes	None	Not reported	Not reported
Montag et al. 2015 [41]	Yes	Yes[43]	Qualitative research and Community outreach – reported in authors’ previous work[43]	Inclusion of pictures and personal stories, emphasize confidentiality, incorporate family and community orientation, information tailored to the local community	Yes Native women clinical staff who had familiarity and experience with the topic and population	Non-Indigenous clinical staff who had familiarity and experience with the topic and population	Participatory research	Not reported

Methodological Quality

The included studies were assessed for their methodological quality from an indigenous perspective using the CREATE tool, and MMAT (Table 4, Table 5 and supplementary file). There was little information reported in the included articles that address the CREATE checklist. Authors of the included studies were contacted to obtain further information and, when provided (n=1), the information was used to appraise the studies. The following standards for measuring quality were applied based on the CREATE tool (Table 4); A study was considered high quality if 10 or more of 14 criteria were met, medium if 6-9 of 14 were met and low if criteria 5 or below were met. Only Hui et al (2021) met the standards for high quality using this tool while the remaining two studies met criteria for low quality evidence [40]. The MMAT assessment (Table 5) returned a quality measurement of 43% (low-medium quality) for Hui et al (2021), 57% for Brown et al (2019) and 71% for Montag et al (2015) (medium quality)[39-41]. This assessment was based on methods used in previous studies which define the quality as low, medium, or high according to the number of ‘Yes’ answers to questions [44, 45]. If less than or equal to 25% of questions were in the affirmative it was a low-quality study, medium if 50% and high if >75% of questions were answered ‘yes’ [44, 45].

Table 4: Quality Assessment -CREATE Tool [31]

Study ID	Q1 *	Q2 *	Q3 *	Q4 *	Q5 *	Q6 *	Q7 *	Q8 *	Q9 *	Q10 *	Q11 *	Q12 *	Q13 *	Q14 *
Hui et al. 2021 [40]	Y	Y	Y	Y	Y	U	U	Y	Y	Y	Y	Y	Y	Y
Brown et al. 2019 [39]	P	U	U	U	U	U	U	U	U	Y	U	Y	U	U
Montag et al. 2015 [41]	Y	N	U	N	N	N	N	N	U	N	U	Y	N	N

Y: Yes, P: Partially, N: No, or U: Unclear

Table 5 Quality Assessment tool MMAT[46]

Citation	Study Design	SQ1	SQ2	Q1	Q2	Q3	Q4	Q5
Hui et al. A, 2021 [40]	Cohort	Y	Y	Y	N	U	U	N
Brown B 2019 [39]	Before-after comparison	Y	Y	N	Y	N	Y	Y
Montag et al. AC 2015 [41]	RCT	Y	Y	Y	Y	N	N	Y

Y: Yes, N: No, or U: Unclear

Discussion

This systematic review demonstrates that there is no evidence of effectiveness of any currently available mHealth intervention designed for indigenous women with young children (0 – 5 years of age) to help assist with behaviour change to healthy choices for themselves and their children. Only three articles found were eligible to be included in this systematic review. These studies aimed to change maternal and child behaviours in the prenatal period and pre-school age group, by provision of a mHealth intervention. The three mHealth interventions that were identified included: a hybrid face-to-face and online prenatal education program; an automated, individualised, three weekly text messages on healthy lifestyle advice for targeting preschool children; and a web-based self-assessment followed by tailored advice related to maternal alcohol drinking. All interventions were applied in a real-life community setting. Statistically significant change was reported in some behaviours/outcomes. However, in the absence of any reported sample size or power calculation; it was not possible to ascertain if significant positive results were due to an inflated Type I error, showing positive results in favour of intervention when in fact the interventions only work due to very large sample sizes.

A recent review on barriers and facilitators of engagement of indigenous peoples with mHealth interventions revealed themes which echo the need for effective and timely co-design practices in this field[47]. Researchers found that barriers to engagement included poor digital design, unreliable technology and internet, repetitive content with limited user input, language barrier, poor cultural representation, non-specific health information and privacy concerns[47]. Co-design has been identified as one of the most important factors determining the uptake of web based therapeutic interventions for indigenous communities/populations[48]. In our review it was deemed important to investigate/ascertain if the digital interventions used were either co-designed in partnership with indigenous communities or were adapted in consultation with the target population to ensure cultural appropriateness; or selected from a pool of previous research/literature on culturally appropriate interventions for the target populations. That information as reported in the included studies is presented in Table 3.

Of the three identified mHealth interventions, two were co-designed with an indigenous population and have shown some positive outcomes in favour of the interventions. Citations of their authors' previous work on co-designing the interventions with indigenous populations were reported for these two interventions in those studies[49, 50]. Although the current evidence on their effectiveness is limited, the authors of these studies are following the iterative process of evidence gathering as suggested in the guideline on the evaluation of complex interventions[51-53]. Therefore, it will be important to see future results on the use of these interventions. Also, due to the diligent process of co-design and insurance of cultural safety, these interventions may provide guidance on the planning and development of work with indigenous populations on provision of maternal behaviour support. The consultation and partnership with Communities reported about the two interventions are in line with the international guidelines on research with indigenous peoples[54].

During the study selection process, eight articles on the development of mHealth interventions for indigenous women were identified that could not be included in this review[23, 24, 55-60]. Upon a further search with the name of the digital interventions reported in the study, no follow up studies on the evaluation of their effectiveness could be retrieved. Further to this, corresponding authors of these articles (n=8) were contacted to inquire of any ongoing studies to evaluate the effectiveness. Except for one, all authors replied to the initial email (n=7/8). Four authors (4/7) mentioned further evaluation work on the identified digital interventions, of which only one has already been published [61]. However, the focus of the publication was not an indigenous population, therefore the study was not included in this review [61]. The other authors (n=3), with ongoing trials, were not able to share the unpublished data. Evidently nearly 40% (3/8) of these mHealth interventions did not evaluate their effectiveness and are no longer active [23, 24, 58]. Although, in response to personal communication, one of the authors [23] mentioned collection of qualitative data on the usage of the app designed for indigenous women [23]; the need for an RCT for the evaluation of effectiveness was also identified. Consistent with the findings of our review, the evidence on the effectiveness of digital health interventions in other populations/settings such as children and adolescents[62], patients with heart failure[63], self-management of chronic obstructive pulmonary disease [64], and sexual health promotion[65] is limited due to the lack of published findings of well-designed and conducted effectiveness evaluation research. There are some health conditions in which the use of mHealth interventions have been well studied, such as asthma[66, 67] and consumption of hazardous amounts of alcohol [68], in which the use of mHealth interventions has been shown to produce positive health outcomes in favour of the interventions[67, 68]. Also, the use of mHealth to promote behavioural change during prenatal care in the general population has been reported to result in significant improvement in behaviour risks, and improved healthy behaviours among pregnant women [69]. A current scoping review on the use and uptake of Web-based therapeutic interventions among indigenous populations across Australia, Canada, New Zealand and USA reported the use of these interventions for conditions such as cardiac care, diabetes, nutrition, mental health, asthma, neonatal care, otitis media, smoking cessation, and substance misuse [70]. For that review, a Web-

based therapeutic intervention was defined as a therapeutic intervention that was self-guided, or a clinician assisted program delivered via the internet to provide guidance, support, and treatment for health conditions. While the evaluation of effectiveness was beyond the scope of that review, it was concluded that these interventions have the potential to improve health and overcome treatment barriers and reduce inequalities for indigenous populations [70].

There are several factors that could lead to the lack of effectiveness evaluation research and in turn scarcity of evidence on any mHealth interventions [71]. Some of these challenges pertain to interventions themselves, whereas some are related to designing a powered RCT with complex interventions which is the common/acceptable approach/method to effectiveness evaluation. Some of the challenges related to mHealth interventions are: rapidly changing technology and the need for intervention to evolve quickly to merely remain functional, let alone upgrade; accessibility of intervention across various operating systems and devices; acceptability and usability of the intervention by target population; real world usage of any intervention or implementation; burden on various stakeholders; adaptation of mHealth interventions to context and population; and their integration into existing healthcare systems. There are also challenges specific to the design of an RCT for mHealth intervention evaluation such as, selecting a suitable context for trial so that the findings are generalisable, choice between conducting a trial with high external versus internal validity, specification, and explanation of the components of the mHealth intervention under evaluation, choice of an appropriate control/comparator group, and data collection methods from mHealth interventions. Given all these, it is arduous to conduct powered RCTs of mHealth interventions [51, 71].

The included studies in this review were appraised for their quality with the CREATE and MMAT[46] tools. The CREATE tool was designed to ensure that the research conducted with indigenous populations is conducted from the perspective of indigenous peoples[31]. There are certain items that authors need to report for the assessors to mark the CREATE tool. While, appraising the included studies in this review, many of those items have not been reported and a further inquiry with the corresponding authors was necessary to access that information. This could have been due to the lack of any guidance on reporting of indigenous research. Recently a guideline was published on reporting of observational studies in indigenous populations[72], however, the studies included in this review were of different methodological design. Another reporting guideline, CONSollDated CritEria or CONSIDER statement, developed by a review of available literature and meta-synthesis, was published in 2019. The guideline provides eight research domains and 17 criteria for the reporting of research involving indigenous peoples. However, except for one, the included studies in the review were published before this date so they cannot be discredited for not following the guideline.

One limitation of this systematic review was single data extraction. Due to the paucity of available information, the data were extracted by one reviewer only. However, once extracted, data were checked for completeness and errors by a second reviewer with more experience in conducting systematic reviews. This limitation is less likely to change the results and conclusions made since the main concern is the lack of relevant literature and research in the field. And the relatively small sample sizes in the included studies (420, 17, 263) meaning the total women included in all studies was 700. Furthermore, the lack of formal power calculations to calculate sample size, to formally assess statistical significance of interventions is problematic. Also, the title and abstract screen for eligibility was performed by one reviewer. This can be perceived as a limitation as there is risk of missing relevant studies as they could be overlooked by single reviewer. We however, kept our decision making more inclusive and the reviewer performing title and abstract screen was instructed to include any study they were unsure about for the full text screening stage. Another potential limitation may be the use of the CREATE tool for risk of bias assessment. The CREATE tool was developed with indigenous and non-indigenous experts and utilises a 14-question checklist which

assesses community engagement, consultation, research governance, and intellectual property to evaluate methodology of studies conducted in the Australian Aboriginal and/or Torres Strait Islander population groups[31]. However, the tool has been widely used in research involving global indigenous populations [32-38]. Therefore, the use of CREATE tool provides additional information about the methodological suitability of the included studies from indigenous viewpoint. Addition of a third aim to the systematic review after the registration of the review protocol may be considered a potential weakness. This was however, not included as a primary aim of the review and did not impact on the conclusion made in the review. Additionally, it was not possible to evaluate differences in the effectiveness of the identified interventions based on participant engagement, intervention nature and context due to the small number of studies included. Also, each study used a different intervention and none of them reported power and sample size calculations. This may be perceived as a weakness of this systematic review. However, it is reflective of the nature of the research and literature available and demonstrates that more research is required in the field.

Conclusion

Overall, the current evidence on the effectiveness of mHealth interventions use with indigenous women is limited. However, there is potential for mHealth to effectively support indigenous women with young children to improve their lifestyle choices for themselves and their children. The interventions that have been codesigned need to have further evaluation research for wider application. New interventions for different contexts and varied indigenous nations are required to be developed, tested for evaluation, and deployed for population use.

[Supplementary file]

Acknowledgements

Authors would like to acknowledge Josepine McGill the research librarian for her assistance with developing the search strategy for this review.

Funding

This systematic review was part of a project funded by the Hospital Research Foundation.

References

1. Truth and Reconciliation Commission of Canada, *Honouring the Truth, Reconciling for the Future*. 2015: Canada. <https://nctr.ca/records/reports/>
2. Axelsson, P., T. Kukutai, and R. Kippen, *The field of Indigenous health and the role of colonisation and history*. *Journal of Population Research*, 2016. **33**(1): p. 1-7. <https://doi.org/10.1007/s12546-016-9163-2>
3. Shah, P.S., et al., *Pregnancy and neonatal outcomes of aboriginal women: a systematic review and meta-analysis*. *Womens Health Issues*, 2011. **21**(1): p. 28-39. 10.1016/j.whi.2010.08.005
4. Akter, S., et al., *Indigenous women's access to maternal healthcare services in lower- and middle-income countries: a systematic integrative review*. *International Journal of Public Health*, 2019. **64**(3): p. 343-353. 10.1007/s00038-019-01200-4
5. Burns, J., C.M. Maling, and N. Thomson, *Summary of Indigenous women's health*. 2010.
6. Waterworth, P., et al., *Factors Influencing the Health Behaviour of Indigenous Australians: Perspectives from Support People*. *PLOS ONE*, 2015. **10**(11): p. e0142323. 10.1371/journal.pone.0142323
7. Batchelor, M., et al., *A Systematic Review of Child Health and Developmental Outcomes Associated with Low Birthweight and/or Small for Gestational Age in Indigenous Children from Australia, Canada and New Zealand*. *International journal of environmental research and public health*, 2021. **18**(23): p. 12669. 10.3390/ijerph182312669
8. Olander, E.K., et al., *Beyond the 'teachable moment' - A conceptual analysis of women's perinatal behaviour change*. *Women and Birth*, 2016. **29**(3): p. e67-e71. 10.1016/j.wombi.2015.10.006
9. NSW Ministry of Health, *The First 2000 Days, Conception to age five, Framework*. 2019: NSW.
10. Ritte, R., et al., *An Australian model of the First 1000 Days: an Indigenous-led process to turn an international initiative into an early-life strategy benefiting indigenous families*. *Glob Health Epidemiol Genom*, 2016. **1**: p. e11. 10.1017/ghg.2016.11
11. Carson, E., et al., *Comparing indigenous mortality across urban, rural and very remote areas: a systematic review and meta-analysis*. *International Health*, 2018. **10**(4): p. 219-227. 10.1093/inthealth/ihy031
12. Australian Institute of Health and Welfare *Aboriginal and Torres Strait Islander Health Performance Framework: summary report 2023*. 2023.
13. Odendaal, W.A., et al., *Health workers' perceptions and experiences of using mHealth technologies to deliver primary healthcare services: a qualitative evidence synthesis*. *Cochrane Database of Systematic Reviews*, 2020(3). 10.1002/14651858.CD011942.pub2
14. Hall, C.S., et al. *Assessing the impact of mHealth interventions in low- and middle-income countries--what has been shown to work?* *Global health action*, 2014. **7**, 25606 DOI: 10.3402/gha.v7.25606.
15. Perkes, S.J., et al., *Aboriginal and Torres Strait Islander Women's Access to and Interest in mHealth: National Web-based Cross-sectional Survey*. *J Med Internet Res*,

2023. **25**: p. e42660. DOI: 10.2196/42660
16. Rice, E.S., et al., *Social media and digital technology use among Indigenous young people in Australia: a literature review*. International Journal for Equity in Health, 2016. **15**(1): p. 81. DOI: 10.1186/s12939-016-0366-0
 17. Kennedy, M., et al., *Codeveloping a multibehavioural mobile phone app to enhance social and emotional well-being and reduce health risks among Aboriginal and Torres Strait Islander women during preconception and pregnancy: A three-phased mixed-methods study*. BMJ Open, 2021. **11**(11). DOI: 10.1136/bmjopen-2021-052545
 18. Reibel, T., et al., *Young Aboriginal women's voices on pregnancy care: Factors encouraging antenatal engagement*. Women and Birth, 2015. **28**(1): p. 47-53. DOI: 10.1016/j.wombi.2014.10.002
 19. Simpson, N., D. Wepa, and K. Bria, *Improving antenatal engagement for Aboriginal women in Australia: A scoping review*. Midwifery, 2020. **91**: p. 102825. DOI: 10.1016/j.midw.2020.102825
 20. Kirmayer, L.J. and G. Brass, *Addressing global health disparities among Indigenous peoples*. The Lancet, 2016. **388**(10040): p. 105-106. DOI: 10.1016/S0140-6736(16)00345-7
 21. Perkes, S.J., et al., *Evaluation of an mHealth Intervention (Growin' Up Healthy Jarjums) Designed With and for Aboriginal and Torres Strait Islander Mothers: Engagement and Acceptability Study*. JMIR Pediatr Parent, 2023. **6**: p. e43673. DOI: 10.2196/43673
 22. Houston, A.G., *Exploring the cultural appropriateness and usefulness of an mHealth program for optimal infant feeding in an urban Aboriginal and Torres Strait Islander health service: growing healthy at Inala*. 2018. accessed via <https://healthbulletin.org.au/articles/exploring-the-cultural-appropriateness-and-usefulness-of-a-mhealth-promotion-program-for-infant-feeding-in-an-urban-aboriginal-health-service-a-qualitative-study/>
 23. Perkes, S.J., et al., *Development of a Maternal and Child mHealth Intervention With Aboriginal and Torres Strait Islander Mothers: Co-design Approach*. JMIR Formative Research, 2022. **6**(7). DOI: 10.2196/36640
 24. Kennedy, M., et al., *Codeveloping a multibehavioural mobile phone app to enhance social and emotional well-being and reduce health risks among Aboriginal and Torres Strait Islander women during preconception and pregnancy: a three-phased mixed-methods study*. BMJ open, 2021. **11**(11): p. e052545. DOI: 10.1136/bmjopen-2021-052545
 25. Chamberlain, C., et al., *Healing The Past By Nurturing The Future: A qualitative systematic review and meta-synthesis of pregnancy, birth and early postpartum experiences and views of parents with a history of childhood maltreatment*. PLOS ONE, 2019. **14**(12): p. e0225441. DOI: 10.1371/journal.pone.0225441
 26. Barrera Jr, M., et al., *Cultural adaptations of behavioral health interventions: a progress report*. Journal of consulting and clinical psychology, 2013. **81**(2): p. 196. DOI: 10.1037/a0027085
 27. Ela, O., Ishaque, S., Bonevski, B., Dowling, A., Brown, C., Hall, K., Canuto, K., Bidargaddi, N., Rissel, C., Briley, A., Roberts, C. , *Digital health interventions for modifying maternal self-care and child-care behaviour amongst Indigenous mothers: a systematic review PROSPERO Registration - CRD42023395710*
 28. Page, M.J., et al., *The PRISMA 2020 statement: an updated guideline for reporting*

- systematic reviews. *BMJ*, 2021. **372**: p. n71. DOI: 10.1136/bmj.n71
29. University of Alberta Library. *Health Sciences Search Filters; Indigenous Peoples*. Available from: <https://guides.library.ualberta.ca/c.php?g=734066&p=5281325>.
 30. Higgins, J.P., et al., *Cochrane handbook for systematic reviews of interventions*. 2019: John Wiley & Sons.
 31. Harfield, S., et al., *Assessing the quality of health research from an Indigenous perspective: the Aboriginal and Torres Strait Islander quality appraisal tool*. *BMC Medical Research Methodology*, 2020. **20**(1): p. 79. DOI: 10.1186/s12874-020-00959-3
 32. Andrews, S., B. Hamilton, and C. Humphreys, *A Global Silence: A Critical Interpretive Synthesis of Aboriginal Mothering Through Domestic and Family Violence*. *Affilia*, 2021. **37**(4): p. 585-604. DOI: 10.1177/08861099211034010
 33. Batchelor, M., et al. *A Systematic Review of Child Health and Developmental Outcomes Associated with Low Birthweight and/or Small for Gestational Age in Indigenous Children from Australia, Canada and New Zealand*. *International Journal of Environmental Research and Public Health*, 2021. **18**, DOI: 10.3390/ijerph182312669.
 34. Bruno, G., et al., *Indigenous Autism in Canada: A Scoping Review*. *Journal of Autism and Developmental Disorders*, 2023. <https://doi.org/10.1007/s10803-023-06045-z>
 35. Davies, A., et al. *Programs Addressing Food Security for First Nations Peoples: A Scoping Review*. *Nutrients*, 2023. **15**, DOI: 10.3390/nu15143127.
 36. Fiolet, R., et al., *Indigenous People's Experiences and Expectations of Health Care Professionals When Accessing Care for Family Violence: A Qualitative Evidence Synthesis*. *Trauma, Violence, & Abuse*, 2020. **23**(2): p. 567-580. DOI: 10.1177/1524838019877998
 37. Povey, J., et al., *An emerging framework for digital mental health design with Indigenous young people: a scoping review of the involvement of Indigenous young people in the design and evaluation of digital mental health interventions*. *Systematic Reviews*, 2023. **12**(1): p. 108. DOI: 10.1186/s13643-023-02215-2
 38. Sanchez, J., et al., *Factors that sustain indigenous youth mentoring programs: a qualitative systematic review*. *BMC Public Health*, 2023. **23**(1): p. 429. DOI: 10.1186/s12889-023-15281-x
 39. Brown, B., et al., *Feasibility of text messaging to promote child health in a rural community on an American Indian reservation*. *Health Education Journal*, 2019. **78**(5): p. 557-569. DOI: 10.1177/0017896919837318
 40. Hui, A., et al., *Impact of remote prenatal education on program participation and breastfeeding of women in rural and remote Indigenous communities*. *EClinicalMedicine*, 2021. **35**: p. 100851. DOI: 10.1016/j.eclinm.2021.100851
 41. Montag, A.C., et al., *Preventing alcohol-exposed pregnancy among an American Indian/Alaska Native population: effect of a screening, brief intervention, and referral to treatment intervention*. *Alcohol Clin Exp Res*, 2015. **39**(1): p. 126-35. DOI: 10.1111/acer.12599
 42. Reid, A.E., et al., *Reducing Barriers for Pregnant Women in Rural/Remote First Nations Communities to Participate in Lifestyle Intervention Program through Community-Oriented Approaches*. *Canadian Journal of Diabetes*, 2012. **36**(5): p. S74-S74. DOI: [10.1016/j.cjcd.2012.07.460](https://doi.org/10.1016/j.cjcd.2012.07.460)
 43. Gorman, J.R., et al., *Creating a culturally appropriate web-based behavioral*

- intervention for American Indian/Alaska Native women in Southern California: the healthy women healthy native nation study.* American Indian Alaska Native Mental Health Research, 2013. DOI: 10.5820/aian.2001.2013.1
44. Badu, E., A.P. O'Brien, and R. Mitchell, *An integrative review of potential enablers and barriers to accessing mental health services in Ghana.* Health Research Policy and Systems, 2018. **16**(1): p. 110. DOI: 10.1186/s12961-018-0380-1
 45. Keelan, K., et al., *Indigenous peoples' experiences and preferences in aged residential care: a systematic review.* AlterNative: An International Journal of Indigenous Peoples, 2021. **17**(2): p. 175-182. DOI: 10.1177/11771801211009167
 46. Hong, Q.N., A. Gonzalez-Reyes, and P. Pluye, *Improving the usefulness of a tool for appraising the quality of qualitative, quantitative and mixed methods studies, the Mixed Methods Appraisal Tool (MMAT).* Journal of Evaluation in Clinical Practice, 2018. **24**(3): p. 459-467. DOI: 10.1111/jep.12884
 47. Whitehead, L., et al., *Barriers to and Facilitators of Digital Health Among Culturally and Linguistically Diverse Populations: Qualitative Systematic Review.* J Med Internet Res, 2023. **25**: p. e42719. DOI: 10.2196/42719
 48. Reilly, R., et al., *Use and uptake of web-based therapeutic interventions amongst Indigenous populations in Australia, New Zealand, the United States of America and Canada: a scoping review.* Syst Rev, 2020. **9**(1): p. 123. DOI: 10.1186/s13643-020-01377-2
 49. Reid, A.E., et al., *Reducing Barriers for Pregnant Women in Rural/Remote First Nations Communities to Participate in Lifestyle Intervention Program through Community-Oriented Approaches.* Canadian Journal of Diabetes, 2012. **36**(5, Supplement): p. S74.
 50. Gorman, J.R., et al., *Creating a culturally appropriate web-based behavioral intervention for American Indian/Alaska Native women in Southern California: the healthy women healthy Native nation study.* Am Indian Alsk Native Ment Health Res, 2013. **20**(1): p. 1-15. DOI: 10.5820/aian.2001.2013.1
 51. Craig, P., et al., *Developing and Evaluating Complex Interventions: The New Medical Research Council Guidance.* BMJ: British Medical Journal, 2008. **337**(7676): p. 979-983. DOI: 10.1136/bmj.a1655
 52. Craig, P. and M. Petticrew, *Developing and evaluating complex interventions: Reflections on the 2008 MRC guidance.* International journal of nursing studies, 2013. **50**(5): p. 585-587. DOI: 10.1016/j.ijnurstu.2012.12.007
 53. Skivington, K., et al., *A new framework for developing and evaluating complex interventions: update of Medical Research Council guidance.* BMJ, 2021. **374**: p. n2061. DOI: 10.1136/bmj.n2061
 54. National Health and Medical Research Council, *Ethical conduct in research with Aboriginal and Torres Strait Islander Peoples and communities: Guidelines for researchers and stakeholders.* 2018, Commonwealth of Australia:: Canberra.
 55. Mairs, R.A., et al., *Gamifying Parenting Education Using an App Developed for Pacific and Other New Zealand Families (Play Kindly): Qualitative Study.* JMIR Serious Games, 2020. **8**(2): p. e15647. DOI: 10.2196/15647
 56. Humphrey, G., et al., *See How They Grow: Testing the feasibility of a mobile app to support parents' understanding of child growth charts.* PLOS ONE, 2021. **16**(2): p. e0246045. 10.1371/journal.pone.0246045
 57. Abbass-Dick, J., et al., *Designing an eHealth Breastfeeding Resource With Indigenous*

- Families Using a Participatory Design*. *J Transcult Nurs*, 2018. **29**(5): p. 480-488. DOI: 10.1177/1043659617753047
58. Hanson, J.D., et al., *Acceptability of an eHealth Intervention to Prevent Alcohol-Exposed Pregnancy Among American Indian/Alaska Native Teens*. *Alcoholism: Clinical and Experimental Research*, 2020. **44**(1): p. 196-202. DOI: 10.1111/acer.14227
59. Dobson, R., et al., *Development of a Culturally Tailored Text Message Maternal Health Program: TextMATCH*. *JMIR Mhealth Uhealth*, 2017. **5**(4): p. e49. DOI: 10.2196/mhealth.7205
60. Ashman, A.M., et al., *Validation of a Smartphone Image-Based Dietary Assessment Method for Pregnant Women*. *Nutrients*, 2017. **9**(1): p. 18. DOI: 10.3390/nu9010018
61. Abbass-Dick, J., et al., *The comparison of access to an eHealth resource to current practice on mother and co-parent teamwork and breastfeeding rates: A randomized controlled trial*. *Midwifery*, 2020. **90**: p. 102812. DOI: 10.1016/j.midw.2020.102812
62. Thabrew, H., et al., *E-Health interventions for anxiety and depression in children and adolescents with long-term physical conditions*. *Cochrane Database of Systematic Reviews*, 2018(8). DOI: 10.1002/14651858.CD012489.pub2
63. Allida, S., et al., *mHealth education interventions in heart failure*. *Cochrane Database of Systematic Reviews*, 2020(7). DOI: 10.1002/14651858.CD011845.pub2
64. Janjua, S., et al., *Digital interventions for the management of chronic obstructive pulmonary disease*. *Cochrane Database of Systematic Reviews*, 2021(4). DOI: 10.1002/14651858.CD013246.pub2
65. Bailey, J., et al., *Computer-based interventions for sexual health promotion: systematic review and meta-analyses*. *International journal of STD & AIDS*, 2012. **23**(6): p. 408-413. DOI: 10.1258/ijsa.2011.011221
66. Ramsey, R.R., et al., *Systematic review of digital interventions for pediatric asthma management*. *The Journal of Allergy and Clinical Immunology: In Practice*, 2020. **8**(4): p. 1284-1293. DOI: 10.1016/j.jaip.2019.12.012
67. Chan, A., et al., *Digital interventions to improve adherence to maintenance medication in asthma*. *Cochrane Database of Systematic Reviews*, 2022(6). DOI: 10.1002/14651858.CD013030.pub3
68. Kaner, E.F., et al., *Personalised digital interventions for reducing hazardous and harmful alcohol consumption in community-dwelling populations*. *Cochrane database of systematic reviews*, 2017(9). DOI: 10.1002/14651858.CD011479.pub2
69. Mauriello, L.M., et al., *Using mHealth to Deliver Behavior Change Interventions Within Prenatal Care at Community Health Centers*. *American Journal of Health Promotion*, 2015. **30**(7): p. 554-562. DOI: 10.4278/ajhp.140515-QUAL-222
70. Reilly, R., et al., *Use and uptake of web-based therapeutic interventions amongst Indigenous populations in Australia, New Zealand, the United States of America and Canada: a scoping review*. *Systematic Reviews*, 2020. **9**(1): p. 123. DOI: 10.1186/s13643-020-01377-2
71. Murray, E., et al., *Evaluating Digital Health Interventions: Key Questions and Approaches*. *American Journal of Preventive Medicine*, 2016. **51**(5): p. 843-851. DOI: 10.1016/j.amepre.2016.06.008
72. Funnell, S., et al., *Improving social justice in observational studies: protocol for the development of a global and Indigenous STROBE-equity reporting guideline*. *International Journal for Equity in Health*, 2023. **22**(1): p. 55. DOI: 10.1186/s12939-023-01893-w

Preprint
JMIR Publications

Supplementary Files

Untitled.

URL: <http://asset.jmir.pub/assets/a4f9b947b89432b374694623ad23a179.docx>

Untitled.

URL: <http://asset.jmir.pub/assets/97012390ed974099d4c3dd2e52225807.docx>

Untitled.

URL: <http://asset.jmir.pub/assets/a9c89c0a4a1407101d8e4f8891d88a51.docx>

CONSORT (or other) checklists

PRISMA Checklist.

URL: <http://asset.jmir.pub/assets/d30cf49d0311d78bd2f66134eabf8e50.pdf>